

5       **EXPRESSION OF G PROTEIN-COUPLED RECEPTORS WITH ALTERED**  
          **LIGAND BINDING AND/OR COUPLING PROPERTIES**

***Related Application***

          This application has been converted from U.S. provisional patent application  
10   Serial No. 60/094,451, filed July 27, 1998, which is incorporated herein, in its entirety,  
      by reference.

***Background of the Invention***

          Cell surface receptors are an important class of proteins involved in cellular  
15   functioning because they are the primary mediators of cell to cell communication. In  
      particular, G protein-coupled receptors (GPCRs) are an important category of cell  
      surface receptors. The medical importance of these receptors is evidenced by the fact  
      that more than 60% of all commercially available prescription drugs work by interacting  
      with known GPCRs.

20       In their resting state, the G proteins, which consist of alpha ( $\alpha$ ), beta ( $\beta$ ) and  
      gamma ( $\gamma$ ) subunits, are complexed with the nucleotide guanosine diphosphate (GDP)  
      and are in contact with the receptors to which they are coupled. When a hormone or  
      other first messenger binds to receptor, the receptor changes conformation and this alters  
      its interaction with the G protein. This spurs the  $\alpha$  subunit to release GDP, and the more  
25   abundant nucleotide guanosine triphosphate (GTP) replaces it, activating the G protein.  
      The G protein then dissociates to separate the  $\alpha$  subunit from the still complexed beta  
      and gamma subunits. Either the  $G\alpha$  subunit, or the  $G\beta\gamma$  complex, depending on the  
      pathway, interacts with an effector. The effector (which is often an enzyme) in turn  
      converts an inactive precursor molecule into an active "second messenger," which may  
30   diffuse through the cytoplasm, triggering a metabolic cascade. After a few seconds, the  
       $G\alpha$  converts the GTP to GDP, thereby inactivating itself. The inactivated  $G\alpha$  may then  
      reassociate with the  $G\beta\gamma$  complex.

          Hundreds, if not thousands, of receptors convey messages through heterotrimeric  
G proteins, of which at least 17 distinct  $G\alpha$  subunit forms have been isolated. Most G  
35   protein-coupled receptors are comprised of a single protein chain that is threaded  
      through the plasma membrane seven times. Such receptors are often referred to as  
      seven-transmembrane domain receptors (STRs). More than a hundred different GPCRs  
      have been found, including many distinct receptors that bind the same ligand, and there  
      are likely many more GPCRs awaiting discovery.

40       The mating factor receptors of yeast cells (STE2 and STE3) span the membrane  
      of the yeast cell seven times and are coupled to yeast G proteins. Heterologous GPCRs

5 can be expressed in yeast cells and can be made to couple to yeast G proteins resulting in the transduction of signals via the endogenous yeast pheromone system signaling pathway which is normally activated by STE2 or STE3. In some cases, such heterologous receptors can be made to couple more effectively to the yeast pheromone system signaling pathway by coexpressing a heterologous G protein  $\alpha$  subunit (e.g. U.S. Patent No. 5,482,835 to King et al), by expressing a chimeric G protein subunit (e.g. 10 WO 94/23025), or by expressing a chimeric G protein-coupled receptor (e.g., U.S. Patent No. 5,576,210 issued to Sledziewski et al.).

The  $\beta\gamma$  subunits of the activated G protein stimulate the downstream elements of the pheromone response pathway, including the Ste20p protein kinase, and a set of 15 kinases that are similar to MEK kinase, MEK (MAP kinase kinase), and MAP kinase of mammalian cells and are encoded by the STE11, STE7, and FUS3/KSS1 genes, respectively (Whiteway et al. 1995. Science. 269:1572).

Members of the family of chemotactic cytokines, which have been proposed to be named "chemokines" for short, are being identified as vital initiators and 20 promulgators of inflammatory and immunological reactions (Oppenheim *et al.* (1991) *Annu Rev Immunol* 9:617). The chemokines range from 8 to 11 kD in MW, are active over a 1 to 100-ng/ml concentration range, and are produced by a wide variety of cell types. They are induced by exogenous irritants and endogenous mediators such as IL-1, TNF, PDGF, and IFN- $\gamma$ . The chemokines bind to specific cell surface receptors with a 25  $K_D$  of 0.4 to 4 nM. These chemokines can be considered "second-order" cytokines that appear to be less pleiotropic than "first-order" proinflammatory cytokines because they are not potent inducers of other cytokines and exhibit more specialized functions in inflammation and repair. As shown in Table 1, some of the chemokines have been assigned to a "chemokine  $\alpha$ " subset based on their gene cluster on chromosome 4 (q12- 30 21) and based on the fact that the first two of their four cysteine groups are separated by one amino acid (C-X-C).

**TABLE 1.** Properties of the chemokine  $\alpha$  subfamily

Cytokine	Cell sources	Exogenous stimulants	Endogenous inducers	Chemotactic or haptotactic responses	Major activities
IL-9	Monocytes	Endotoxin	IL-1	Neutrophils	Activates PMN

	Neutrophils Fibroblasts Endothelial cells Keratinocytes Large granular lymphs T lymphocytes	Mitogens Particulates Viruses	TNF IFN- $\gamma$ costimulant IL-3	Basophils Unstimulated T cells Melanoma Cells	$\uparrow$ Neutrophil adhesion $\downarrow$ Basophil histamine $\uparrow$ Keratinocyte growth Acute Inflammation
GRO- $\alpha\beta\gamma$ /mu/KC muMIP-2 $\alpha$ $\beta$	Monocytes Fibroblasts Endothelial cells	Endotoxin	IL-1 TNF	Neutrophils	Degranulates PMN $\uparrow$ Melanoma cell growth $\uparrow$ Fibroblast growth Acute inflammation
CTAP III/ $\beta$ TG $\beta$ TG/NAP-2 PF-4	Monocytes Platelets Monocytes Platelets Platelets	Platelet activators Platelet activators Platelet activators		Fibroblasts Neutrophils Fibroblasts	$\uparrow$ Fibroblast growth Activates PMN $\uparrow$ Fibroblast growth Reverses immune suppression $\uparrow$ I-CAM1 on E.C. $\uparrow$ Chronic inflammation
IP- 10/muCCRG -2	Monocytes Fibroblasts Endothelial cells Keratinocytes	Endotoxin	IFN- $\gamma$	Monocytes Activated T lymphocytes	
ENA-78	Epithelial cells		IL-1 TNF	Neutrophils	Activates PMN

- 5 This chemokine  $\alpha$  group includes IL-8, melanoma growth-stimulating activity (MGSA/GRO), platelet factor 4 (PF-4),  $\beta$  thromboglobulin ( $\beta$ TG), IP-10, and ENA-78.

IL-8 is produced by many cell types including NK cells and T lymphocytes in response to exogenous stimuli such as polyclonal mitogens, injurious stimuli, and infectious agents, as well as proinflammatory cytokines such IL-1 and TNF. IL-8 is a  
10 chemoattractant of neutrophils, basophils, and a small proportion (10% or less) of resting CD4<sup>+</sup> and CD8<sup>+</sup> lymphocytes. IL-8 additionally activates neutrophil enzyme release. IL-8 is also haptotactic for melanocytes and is a comitogenic stimulant of keratinocytes.

IL-8 promotes the adherence of neutrophils to endothelial cells. IL-8 does so by  
15 inducing neutrophils to express  $\beta_2$  integrins. Neutrophils then extravasate by moving between the endothelial cell junctions and through the basement membrane to

5 accumulate in the tissues. Intracutaneous injections of IL-8 cause a rapid local  
neutrophilic infiltration peaking with 3 hours. Intravenous administration of IL-8 does  
not induce systemic sequelae of elevation of acute-phase proteins or fever but does  
induce a neutrophilia. Intravenous administration of IL-8 also specifically reduces local  
peripheral inflammatory responses to IL-8, fMLP, and C5a (Hechman *et al.* (1990)  
10 *FASEB J* 4:890). This transient anti-inflammatory effect of IL-8 probably can be  
attributed to desensitization of neutrophils by systemically distributed IL-8.

Two distinct but homologous (70% at the amino acid level) receptors for IL-8  
have been cloned. The IL-8 receptors are members of the rhodopsin receptor family and  
have a seven transmembrane spanning region (Holmes *et al.* (1991) *Science* 253:1278;  
15 Murphy *et al.* (1991) *Science* 253:1280). The receptors are probably coupled to G-  
proteins, transduce phosphoinositol hydrolysis, and are capable of rapid elevation of  
diacylglycerol and cytosolic Ca<sup>2+</sup> levels, which may lead to activation of protein kinase  
C (Thelen *et al.* (1988) *FASEB J* 2:2702). IL-8 receptors are expressed by neutrophils,  
which display both types of IL-8R and their expression is unregulated by G-CSF (A.  
20 Lloyd *et al.*, *unpublished results*). Mature neutrophils express about 20,000 receptors  
per cell. Myelocytic lines and basophils express several thousand receptors per cell.

IL-8 like molecules have been identified in rabbits, sheep, and other species. IL-  
8 is in the circulation of patients with systemic inflammatory reactions or severe trauma.  
IL-9 has readily been detected in inflammatory sites such as in the synovial fluid in  
25 rheumatoid arthritis (Brennan *et al.* (1990) *Eur J. Immunol* 20:2141), extracts of  
psoriatic skin (Schroder *et al.* (1986) *J Invest Dermatol* 87:53, and in the circulation of  
patients in septic shock (Van Zee *et al.* (1991) *J Immunol* 146:3478-3482). Thus IL-8 is  
implicated as a major participant in acute as well as more prolonged inflammatory  
reactions.

30 MGSA, as its name implies, was first discovered as a factor that accelerated the  
growth of melanoma cell lines and also as a product of oncogene transfected cell lines  
(GRO). MGSA/GRO competes for the type II, but not type I, IL-8 receptor on  
myelocytic cells (Lee (1992) *J Biol Chem* 267:16283-16287) and is also a potent  
chemoattractant, as well as activator of neutrophils. MGSA as well as IL-8 has been  
35 extracted from psoriatic tissues (Sticherling *et al.* (1991) *J Invest Dermatol* 96:26).

GRO has three variants ( $\alpha$ ,  $\beta$ , and  $\gamma$ ), which exhibit about 95% homology in their  
amino acid sequences. They are probably homologues of murine macrophage derived  
KC, macrophage derived inflammatory peptides MIP-2 $\alpha$  and MIP-2 $\beta$ . Murine MIP-2 $\alpha$   
and MIP-2 $\beta$  both compete with equal affinity for type II receptors for IL-8 and  
40 chemoattract human as well as murine neutrophils (Lee *et al.* (1992) *J. Biol Chem*  
267:16283-16287). MIP-2 is also reported to degranulate murine neutrophils, resulting

5 in the release of lysosomal enzymes. Local *in vivo* injections of MIP-2 results in neutrophil accumulation and MIP-2 has been isolated from sites of wound healing. MIL-2 is a costimulator of hematopoietic colony formation by CSF-1 and GM-CSF, but the *in vivo* relevance of this observation remains to be established. It is most likely that GRO/MIP-2 inflammatory activities overlap considerably with those of IL-8, and GRO  
10 is therefore probably also a major inflammatory mediator.

*PF-4* and *CTAP III*, the precursor of  $\beta$ TG, are both present in platelet granules and are released by inducers of platelet aggregation. Consequently, they become available at sites of injury, hemorrhage, and thromboses. Both are reported to chemoattract and to stimulate fibroblasts, presumably for repair purposes. In addition, a  
15 70 amino acid breakdown product of  $\beta$ TG known as neutrophil attracted peptide 2 (*NAP-2*) is a chemoattractant and activator of neutrophils, albeit at 100-fold higher concentrations than IL-8. *NAP-2* also competes for the type II IL-8 receptor with about one-hundredth of the affinity of IL-8 (Leonard *et al.* (1991) *J. Leukoc Biol* **49**:258). However, since at the site of platelet aggregation, high levels of *NAP-2* can be released,  
20 it is thought to be an active participant in attracting inflammatory cells to such sites.

*ENA-78* is the most recently cloned member of the chemokine  $\alpha$  subfamily (Walz *et al.* (1991) *J Exp Med* **174**:1355). *ENA-78* is produced by an epithelial cell line in response to IL-1 and TNF. In cross-desensitization experiments, *ENA-78* also utilizes the type II receptor for IL-8 and GRO and is a chemoattractant and activator of  
25 neutrophils.

*IP-10* is produced by macrophages, endothelial cells, and keratinocytes in response to IFN- $\gamma$ . The pathophysiological functions of *IP-10* remain unclear, but antibodies to *IP-10* react with many cell types present at sites of delayed hypersensitivity reactions and *IP-10* has been extracted from psoriatic plaques (Gotlieb *et al.* (1988) *J*  
30 *Exp Med* **168**:941). Thus *IP-10* can presumably be produced by many cell types and probably participates in chronic inflammation and delayed hypersensitivity responses. A stable recombinant human *IP-10* was recently produced by Dr. K. Matsushima (*personal communication*). We have shown that this rh*IP-10* is a moderately potent *in vitro* chemoattractant of human monocytes, but not neutrophils. In addition, this *IP-10*  
35 also is a moderately potent chemoattractant for previously activated CD4 and CD8 T lymphocytes and promotes adhesion of lymphocytes to endothelial cells (D. Taub *et al.*, *unpublished results*). These observations predict that *IP-10* will probably be a participant in chronic cell-mediated inflammatory reactions.

There are two known human subtypes of the IL-8 receptor. Subtype A(CXCR1)  
40 binds IL-8 with high affinity ( $K_d=0.1$ nM) but shows very low affinity binding to GRO $\alpha$

5 and NAP-2 ( $K_d > 100\text{nM}$ ). Subtype B (CXCR2) binds all three ligands with high affinity.

Overall, the chemokine family members appear to be very potent and pivotal chemoattractants and activators of inflammatory cells and fibroblasts. The  
10 aforementioned inflammatory cytokines have myriad effects on cell growth and differentiation. Some of these effects are indirect and based on the capacity of cytokines to induce the production of a cascade of other cytokines. Therapeutic usefulness of cytokines and their inhibitors is growing and should accelerate.

In recent years drug discovery has been advanced by expression of heterologous receptors in living cells. However, due to the complexities inherent in such  
15 heterologous expression studies, the development of reliable assays to search for modulators of these receptors has presented particular challenges. For example, it is often difficult to obtain sufficient expression of heterologous G protein-coupled receptors or to achieve functional integration of the G protein-coupled receptor into a signaling pathway. Price et al. have reported the functional coupling of a rat A2a  
20 adenosine receptor into the yeast pheromone response pathway (Price et al. (1996) *Molec. Pharmacol.* 50:829-837), but only achieved effective coupling when they coexpressed the native yeast G protein GPA1, expressed from a plasmid, in the yeast cells. They were unable to achieve effective coupling using either a mammalian  $G\alpha$  subunit protein or a chimeric  $G\alpha$  subunit protein expressed in the yeast cells. The  
25 development of expression and coupling systems for expression of modified G protein coupled receptors which display altered ligand binding and/or coupling in host cells would be of tremendous benefit in the development of improved drug screening assays for modulators of G protein coupled receptors.

### 30 ***Summary of the Invention***

The present invention is based, in part, upon studies investigating the coupling of the IL8A receptor to the pheromone response pathway in yeast through a Gpal-G $\alpha$ i2 chimera. By random mutagenesis of this receptor, mutants have been identified that show altered coupling behavior to the ligands GRO $\alpha$  and NAP-2 as well as showing an  
35 enhanced response to IL8. Based upon these studies, an amino acid motif has been identified in the IL8 receptor. Mutations made in this region can affect ligand binding and/or coupling of the receptor to the signal transduction machinery of the cell. Other receptors also have this motif and corresponding mutations can be made in such receptors, e.g., the galanin receptor-1 to enhance signaling by those receptors through  
40 Gpal-G $\alpha$ i2.

5           Thus, the present invention provides host cells that express modified G protein coupled receptors which are functionally integrated into the pheromone response pathway of the cells such that the cells display ligand dependent responses to the modified receptor which differ from responses seen upon modulation of the wild type receptor. The cells of the invention can be used to identify modulators (*e.g.*, agonists or  
10   antagonists) of such receptors. In particular, the invention provides cells expressing modified mammalian IL-8 or galanin receptors (*e.g.*, IL8A or galanin-1 receptors). When yeast cells are used as the host cells, yeast cells may achieve functional coupling of the heterologous G protein coupled receptor using a mammalian G protein subunit protein or a chimeric G protein subunit expressed in the yeast cells. The invention also  
15   provides for mammalian cells which express these modified forms of G protein-coupled receptors.

          In one embodiment, the invention provides a mutant mammalian G-protein coupled receptor having a sequence which varies from a wild type G protein-coupled receptor having a wild type amino acid sequence comprising an amino acid motif  
20   [X<sub>1</sub>X<sub>2</sub>X<sub>3</sub>X<sub>4</sub>] lying near the carboxy terminal end of said domain, wherein:  
                  X<sub>1</sub> denotes an amino acid residue at position 1 of said motif and is selected from the group consisting of Phe, Leu, Val, and Tyr;  
                  X<sub>2</sub> denotes an amino acid residue at position 2 of said motif and is selected from the group consisting of Phe, Lys and Gln;  
25                X<sub>3</sub> denotes an amino acid residue at position 3 of said motif and is selected from the group consisting of Leu, Arg, Glu, Asn, Gln, Ser, Ala, Leu ; and  
                  X<sub>4</sub> denotes an amino acid residue at position 4 of said motif and is selected from the group consisting of Ala, Cys, Asp, Glu, Gly, Ser, Thr and Tyr; and  
                  wherein said mutant receptor comprises a seventh transmembrane domain  
30   with a carboxy terminal end;  
                  at least one point mutation at a position in said amino acid motif; such that upon interaction with a ligand to modulate a signal transduction pathway in a cell, a signal generated by said mutant receptor is greater than a signal generated upon interaction of said ligand with a wild type G protein-coupled receptor.

35           In one embodiment, the host cell is a yeast cell. In a preferred embodiment, the receptor acts as a surrogate for an endogenous yeast pheromone receptor in a pheromone response pathway of said cell. In a particularly preferred embodiment, the cell belongs to the species *Saccharomyces cerevisiae*.

          In one embodiment, the cell is a mammalian cell.

40           In one embodiment, interaction of ligand with a wild type G protein-coupled receptor generates no detectable signal. In one embodiment, the mutant receptor comprises an Arg or Lys at position 4.

5 In one embodiment, the heterologous receptor is an IL8A receptor.

In a preferred embodiment the point mutation in the receptor is selected from the group consisting of : Arg to Trp at position 73, Met to Ile at position 246; and Gly to Arg at position 320.

10 In one embodiment, the ligand of the receptor is interleukin 8 (IL8) or melanoma growth-stimulating activity-alpha (MGSA/GRO $\alpha$ ).

In a preferred embodiment, the receptor is a human receptor.

In another preferred embodiment the receptor is selected from the group consisting of human galanin-1 receptor, somatastatin receptor type I, somatastatin receptor type II, somatastatin receptor type III, and human nociceptin receptor. In  
15 particularly preferred embodiments, the receptor is a human galanin-1 receptor.

In preferred embodiments, the receptor comprises an amino acid sequence LAYSNSSVNPIIYAFLSEN[FRKR]YKQV (SEQ ID NO: 1) wherein said mutant amino acid motif within said sequence is [FRKR] (SEQ ID NO: 2).

In another aspect, the invention provides a recombinant yeast cell having a  
20 mutant form of a mammalian G protein-coupled receptor expressed in a membrane of said yeast cell, said mutant receptor having a sequence that varies from a wild type G protein-coupled receptor having a wild type amino acid sequence comprising an amino acid motif [X<sub>1</sub>X<sub>2</sub>X<sub>3</sub>X<sub>4</sub>] lying near the carboxy terminal end of said domain, wherein:

X<sub>1</sub> denotes an amino acid residue at position 1 of said motif and  
25 is selected from the group consisting of Phe, Leu, Val, and Tyr;

X<sub>2</sub> denotes an amino acid residue at position 2 of said motif and is selected from the group consisting of Phe, Lys and Gln;

X<sub>3</sub> denotes an amino acid residue at position 3 of said motif and is selected from the group consisting of Leu, Arg, Glu, Asn, Gln, Ser, Ala, Leu ; and

30 X<sub>4</sub> denotes an amino acid residue at position 4 of said motif and is selected from the group consisting of Ala, Cys, Asp, Glu, Gly, Ser, Thr and Tyr; and wherein said mutant receptor comprises:

a seventh transmembrane domain with a carboxy terminal end;  
and

35 at least one point mutation at a position on said amino acid motif; wherein said mutant receptor acts as a surrogate for an endogenous yeast pheromone receptor in a pheromone response pathway of the yeast cell, such that upon interaction with a ligand to modulate signal transduction in said pheromone response pathway, a signal generated by said mutant receptor is greater than a signal generated upon  
40 interaction of said ligand with a wild type G protein-coupled receptor.

In a preferred embodiment, the yeast cell comprises a human receptor.



5 In another preferred embodiment, the yeast cell expresses a heterologous receptor selected from the group consisting of human galanin-1 receptor, somatastatin receptor type I, somatastatin receptor type II, somatastatin receptor type III, and human nociceptin receptor. In another preferred embodiment, the yeast cell expresses an IL8A receptor.

10 In another preferred embodiment, the yeast cell expresses a mammalian, chimeric, and/or mutant G protein subunit. In another preferred embodiment, the yeast cell expresses a mammalian, chimeric, and/or mutant  $G\alpha$  subunit. In a preferred embodiment, the yeast cell expresses a GPA41- $G\alpha_{i2}$  subunit. In another embodiment, the yeast cell expresses a GPA41- $G\alpha_{i1}$  or GPA41- $G\alpha_{i3}$  subunit.

15 In another embodiment, the yeast cell expresses a STE18- $G\gamma 2$  subunit.

In another embodiment, the yeast cell expresses a mammalian  $G_{\alpha S}$ E10K subunit. In another embodiment, the yeast cell expresses a mammalian  $G_{\alpha S}$ D229S subunit or a mammalian  $G_{\alpha S}$ E10K + D229S subunit.

20 In one embodiment, the yeast cell is a mutant cell having a pheromone response pathway that is desensitized at slower rate than a wild type strain under the same conditions of continuous stimulation of the pheromone response pathway. In one embodiment, the yeast cell has a *ste14* mutation.

25 In another embodiment, the yeast cell of claim 15, which has a *ste2* or *ste3* mutation. In another embodiment, the endogenous pheromone gene is not functionally expressed in the yeast cell. In yet another embodiment, an endogenous *FAR1* gene is not functionally expressed in the yeast cell. In a further embodiment, an endogenous *SST2* gene is not functionally expressed in the yeast cell.

In another embodiment, the yeast cell comprises a detectable or selectable marker that is activated by a pheromone response pathway of the yeast cell.

30 In one embodiment, the selectable marker comprises a pheromone-responsive promoter operably linked to a selectable gene.

In one embodiment, the pheromone-responsive promoter is the *FUS1* promoter. In another embodiment, the marker gene is a HIS 3 gene or a LacZ gene.

35 In another embodiment, the yeast cell further comprises a heterologous polypeptide, wherein the heterologous polypeptide is transported to a location allowing interaction with the region of said receptor expressed in the cell membrane and wherein modulation of the signal transduction of said receptor by the heterologous polypeptide provides a detectable signal.

40 In one embodiment, the heterologous polypeptide includes a signal sequence. In another embodiment, the signal sequence corresponds to a leader peptide of the *Saccharomyces cerevisiae*  $\alpha$  factor or a-factor.

In preferred embodiments, the yeast cells are *Saccharomyces cerevisiae* cells.

5 In another aspect, the invention pertains to a method for identifying a modulator of a mammalian G-coupled protein receptor expressed by a yeast cell, comprising:

- (i) contacting a mixture of yeast cells with a test compound;
- (ii) allowing cells within the mixture to generate a detectable signal; and
- (iii) identifying the test compound as a modulator of said receptor expressed by

10 the yeast cell.

### *Description of the Drawings*

15 Figure 1 shows ligand induced growth of cells expressing wild type or mutant IL8 receptors and the indicated ligand.

Figure 2 shows amino acid sequence of the rabbit IL8A receptor showing putative  
20 membrane spanning domains. Arg73 (1st intracellular loop), Met246 (3rd intracellular loop) and Gly320 (C-terminal tail) are marked in red.

Figure 3 shows enhanced growth phenotype of mutant receptors.

25 Figure 4 shows FUS1-lacZ activity in autocrine strains expressing wild type or mutant receptors.

Figure 5 shows activation of wild type or mutant IL8 receptors by exogenous IL8 (A), GRO $\alpha$  (B) or NAP-2 (C).

30 Figure 6 shows saturation binding of [ $^{125}$ I] IL8 to membranes from yeast cells expressing the wild type IL8 receptor or the Arg73Trp Gly320Arg double mutant.

Figure 7 shows ligand dependent growth of yeast cells expressing wild type or mutant  
35 human Galanin receptor.

Figure 8 shows Ca $^{2+}$  flux in mammalian HEK293/G $\alpha$ 16 cells expressing wild type or mutant IL8 receptors.

40 Figure 9 shows pertussis toxin sensitivity of the ligand induced Ca $^{2+}$  flux in mammalian HEK293/G $\alpha$ 16 cells expressing wild type or mutant IL8 receptors

5

### ***Detailed Description of the Invention***

10 The present invention provides modified forms of G protein coupled receptors which display altered ligand binding and/or coupling properties. The invention also provides cells expressing such receptors and assays utilizing these cells for screening and identifying pharmaceutically effective compounds that specifically modulate the activity of a these modified forms of G protein coupled receptors. Yeast or mammalian cells can be used to express such receptors. The subject assays enable rapid screening of  
15 large numbers of compounds (e.g., compounds in a library) to identify those which are receptor agonists or antagonists. Compositions of matter, such as these novel receptors, novel recombinant yeast cells and novel gene constructs, are also embraced by the present invention. The instant assays provide a convenient format for discovering compounds which can be useful in modulating cellular function, as well as in  
20 understanding the pharmacology of compounds that specifically interact with these modified forms G protein coupled receptors.

In the practice of the instant invention, standard techniques known in the art can be used. See for example, Sherman. 1991. *Methods Enzymol.* 194:3; Sherman and Hicks. 1991. *Methods Enzymol.* 194:21; Sambrook et al. *Molecular Cloning A*  
25 *Laboratory Manual*, 2nd Ed., ed. by Sambrook, Fritsch and Maniatis (Cold Spring Harbor Laboratory Press: 1989 or 1991 edition.

Before further description of the invention, certain terms employed in the specification, examples and appended claims are, for convenience, collected here.

#### 30 **I. Definitions**

The term "compound" as used herein (e.g., as in "test compound") is meant to include both exogenously added test compounds and peptides endogenously expressed by a cell, e.g., from a peptide library. For example, in certain embodiments, the reagent cell also produces the test compound which is being screened. For instance, the reagent  
35 cell can produce. e.g., a test polypeptide, a test nucleic acid and/or a test carbohydrate which is screened for its ability to modulate the heterologous receptor activity. In such embodiments, a culture of such reagent cells will collectively provide a library of potential effector molecules and those members of the library which either agonize or antagonize the receptor function can be selected and identified. Moreover, it will be  
40 apparent that the reagent cell can be used to detect agents which transduce a signal via the receptor of interest.

5           In one embodiment, the compound is a ligand for the receptor, e.g., a compound which interacts with the receptor, whether naturally occurring or nonnaturally occurring. As used herein, the language "interacts with" includes receptor-ligand interactions, e.g., binding interactions. Preferably, when a ligand "interacts with" a receptor, the receptor transduces a signal in the cell expressing the receptor, such that a signal transduction  
10       pathway is modulated, e.g., affected in the cell.

          In other embodiments, the test compound is exogenously added. In such embodiments the test compound is contacted with the reagent cell. Exemplary compounds which can be screened for activity include, but are not limited to, peptides, nucleic acids, carbohydrates, small organic molecules, and natural product extract  
15       libraries. In such embodiments, both compounds which agonize or antagonize the receptor mediated signaling function can be selected and identified.

          The term "non-peptidic compound" is intended to encompass compounds that are comprised, at least in part, of molecular structures different from naturally-occurring L-amino acid residues linked by natural peptide bonds. However, "non-peptidic  
20       compounds" are intended to include compounds composed, in whole or in part, of peptidomimetic structures, such as D-amino acids, non-naturally-occurring L-amino acids, modified peptide backbones and the like, as well as compounds that are composed, in whole or in part, of molecular structures unrelated to naturally-occurring L-amino acid residues linked by natural peptide bonds. "Non-peptidic compounds" also  
25       are intended to include natural products.

          As used herein, "recombinant cells" include any cells that have been modified by the introduction of heterologous DNA. Exemplary control cells include cells that are substantially identical to the recombinant cells, but do not express one or more of the proteins encoded by the heterologous DNA, e.g., do not include or express a reporter  
30       gene construct, receptor or test polypeptide, or express a different heterologous DNA (e.g., a cell that expresses a different GPCR that couples to the same G protein as that of the GPCR whose activity is being examined).

          As used herein, "heterologous DNA" or "heterologous nucleic acid" includes DNA that does not occur naturally as part of the genome in which it is present or which  
35       is found in a location or locations in the genome that differs from that in which it occurs in nature or which is operatively linked to DNA to which it is not normally linked in nature (i.e., a gene that has been operatively linked to a heterologous promoter). Heterologous DNA is not naturally occurring in that position or is not endogenous to the cell into which it is introduced, but has been obtained from another cell. Heterologous  
40       DNA can be from the same species or from a different species. In some embodiments, it is mammalian, e.g., human. Any DNA that one of skill in the art would recognize or

5 consider as heterologous or foreign to the cell in which is expressed is herein  
encompassed by the term heterologous DNA. Examples of heterologous DNA include,  
but are not limited to, genes which encode proteins that amplify signals transduced via  
the pheromone response pathway, DNA that encodes test polypeptides, receptors,  
reporter genes, transcriptional and translational regulatory sequences, or selectable or  
10 traceable marker proteins, such as a protein that confers drug resistance.

The terms "heterologous protein", "recombinant protein", and "exogenous  
protein" are used interchangeably throughout the specification and refer to a polypeptide  
which is produced by recombinant DNA techniques, wherein generally, DNA encoding  
the polypeptide is inserted into a suitable expression vector which is in turn used to  
15 transform a host cell to produce the heterologous protein. That is, the polypeptide is  
expressed from a heterologous nucleic acid.

As used herein, the term "high copy number plasmid" refers to a plasmid which  
exists in at least 5, or more, copies per cell, and more preferably in at least 10-20 copies  
per cell. The term "low copy number plasmid" refers to a plasmid which exists in fewer  
20 than 5 copies per cell, more preferably 2-3 copies, or less, per cell.

As used herein, the term "extracellular signal" is intended to encompass  
molecules and changes in the environment that are transduced intracellularly via cell  
surface proteins that interact, directly or indirectly, with the extracellular signal. An  
extracellular signal or effector molecule includes any compound or substance that in  
25 some manner alters the activity of a cell surface protein. Examples of such signals  
include, but are not limited to, molecules such as acetylcholine, growth factors and  
hormones, lipids, sugars and nucleotides that bind to cell surface receptors and modulate  
the activity of such receptors. The term, "extracellular signal" also includes as yet  
unidentified substances that modulate the activity of a cellular receptor, and thereby  
30 influence intracellular functions. Such extracellular signals are potential  
pharmacological agents that may be used to treat specific diseases by modulating the  
activity of specific cell surface receptors.

As used herein, "heterologous G protein receptor" (e.g., a heterologous adenosine  
receptor) is encoded by heterologous DNA and, upon expression of this heterologous  
35 DNA in a recombinant cell, the heterologous receptor is expressed in the recombinant  
cell.

The term "signal transduction" is intended to encompass the processing of  
physical or chemical signals from the extracellular environment through the cell  
membrane and into the cell, and may occur through one or more of several mechanisms,  
40 such as activation/inactivation of enzymes (such as proteases, or other enzymes which  
may alter phosphorylation patterns or other post-translational modifications), activation

5 of ion channels or intracellular ion stores, effector enzyme activation via guanine nucleotide binding protein intermediates, formation of inositol phosphate, activation or inactivation of adenylyl cyclase, direct activation (or inhibition) of a transcriptional factor and/or activation.

10 A "signaling pathway" refers to the components involved in "signal transduction" of a particular signal into a cell. The term "endogenous signaling pathway" indicates that some or all of the components of the signaling pathway are naturally-occurring components of the cell. An example of such a pathway is the endogenous pheromone response pathway of yeast.

15 The term "functionally couples to" (as in a receptor that is "functionally integrated into a signaling pathway in a cell" or "functionally integrated into an endogenous yeast signaling pathway" or "functionally expressed by a host cell") is intended to refer to the ability of the receptor to be expressed at the surface of the cell and the ability of the expressed receptor to bind to modulators (e.g., a ligand of the receptor) and transduce signals into the cell via components of a signaling pathway of  
20 the cell. For example, a G protein-coupled receptor (GPCR) which is functionally integrated into an endogenous pheromone response or signaling pathway of a yeast cell is expressed on the surface of the yeast cell, couples to a G protein within the yeast cell and transduces a signal in that yeast cell upon binding of a modulator to the receptor. For example, a G protein subunit, e.g., a chimeric, mutant or heterologous subunit, that  
25 is functionally integrated into a yeast cell may be capable of coupling both to the GPCR and to the other G protein subunits, which can also be endogenous to the yeast cell, can be chimeric, or can be heterologous. Alternatively, the G protein subunit can be constitutively active such that it need not be coupled to a heterologous GPCR. A transduced signal may be detected by measuring any one of a number of responses to  
30 mating factors which occur in a yeast cell, e.g., growth arrest or transcription of an indicator gene responsive to signals produced by modulation of a pheromone response pathway or any biochemical changes.

35 The term "indicator gene" generically refers to an expressible (e.g., able to transcribed and (optionally) translated) DNA sequence which is expressed in response to a signal transduction pathway modulated by a target receptor or ion channel. Exemplary indicator genes include unmodified endogenous genes of the host cell, modified endogenous genes, or a reporter gene of a heterologous construct, e.g., as part of a reporter gene construct.

40 The term "endogenous gene" is intended to refer to a gene in a cell that is naturally part of the genome of the cell and which, most preferably, is present in its natural location in the genome (as opposed to "heterologous" DNA which has been

5 introduced into the cell). Likewise, the term "endogenous protein" is intended to include proteins of a cell that are encoded by endogenous genes of the cell.

An endogenous gene that is to be used as an indicator gene may comprise the natural regulatory elements of the gene (e.g., the native promoter/enhancer elements that naturally regulate expression of the gene) or the endogenous gene can be "operatively  
10 linked to" (i.e., functionally coupled to) a "heterologous promoter" (or other heterologous regulatory elements). A "heterologous promoter" refers to a promoter that does not naturally regulate the gene to which the heterologous promoter is operatively linked. For example, an endogenous yeast gene that is not normally pheromone-responsive can be operatively linked to a heterologous promoter that is responsive to  
15 signals produced by the yeast pheromone system to thereby confer pheromone responsiveness on the endogenous yeast gene. Methods of using endogenous yeast genes as indicator genes are described further in PCT Publication WO 98/13513, the contents of which are hereby expressly incorporated herein by this reference.

The term "detecting an alteration in a signal produced by an endogenous  
20 signaling pathway" (e.g., an endogenous yeast signaling pathway) is intended to encompass the detection of alterations in endogenous second messengers produced upon activation of components of the endogenous signaling pathway, alterations in endogenous gene transcription induced upon activation of components of the endogenous signaling pathway, and/or alterations in the activity of an endogenous  
25 protein(s) upon activation of components of the endogenous signaling pathway. In certain embodiments, the term "detecting an alteration in a signal produced by an endogenous signaling pathway" can also encompass assaying general, global changes to the cell such as changes in cell growth or cell morphology.

As used herein, a "reporter gene construct" refers to a nucleic acid that includes a  
30 "reporter gene" operatively linked to a transcriptional regulatory sequences. Transcription of the reporter gene is controlled by these sequences. The activity of at least one or more of these control sequences is directly or indirectly regulated by the target receptor protein. The transcriptional regulatory sequences include the promoter and other regulatory regions, such as enhancer sequences, that modulate the activity of  
35 the promoter, or regulatory sequences that modulate the activity or efficiency of the RNA polymerase that recognizes the promoter, or regulatory sequences which are recognized by effector molecules, including those that are specifically induced by interaction of an extracellular signal with the target receptor. For example, modulation of the activity of the promoter may be effected by altering the RNA polymerase binding  
40 to the promoter region, or, alternatively, by interfering with initiation of transcription or elongation of the mRNA. Such sequences are herein collectively referred to as

5 transcriptional regulatory elements or sequences. In addition, the construct may include sequences of nucleotides that alter translation of the resulting mRNA, thereby altering the amount of reporter gene product. The reporter gene constructs of the present invention provide a detectable readout in response to signals transduced in response to modulation of a heterologously expressed receptor.

10 The term "modulation", as in "modulation of a (heterologous) G protein coupled receptor" and "modulation of a signal transduction activity of a receptor protein" is intended to encompass, in its various grammatical forms, induction and/or potentiation, as well as inhibition and/or downregulation of receptor activity and/or one or more signal transduction pathways downstream of a receptor.

15 Agonists and antagonists are "receptor effector" molecules that modulate signal transduction via a receptor. Receptor effector molecules are capable of binding to the receptor, though not necessarily at the binding site of the natural ligand or otherwise modulating the activity of the receptor, for example, by influencing the activity of components which regulate the receptor, or which function in the signal transduction  
20 initiated by the receptor. Receptor effectors can modulate signal transduction when used alone, i.e. can be surrogate ligands, or can alter signal transduction in the presence of the natural ligand or other known activators, either to enhance or inhibit signaling by the natural ligand. For example, "antagonists" are molecules that block or decrease the signal transduction activity of receptor, e.g., they can competitively, noncompetitively,  
25 and/or allosterically inhibit signal transduction from the receptor, whereas "agonists" potentiate, induce or otherwise enhance the signal transduction activity of a receptor.

The term "surrogate ligand" refers to an agonist which induces signal transduction from a receptor; the agonist is a surrogate in that it is not the natural ligand of the receptor.

30 The term "autocrine cell", as used herein, refers to a cell which produces a substance which can stimulate a receptor located on or within the same cell as that which produces the substance. For example, wild-type yeast MAT $\alpha$  and MATa cells are not autocrine. However, a yeast cell which produces both  $\alpha$ -factor and  $\alpha$ -factor receptor, or both a-factor and a-factor receptor, in functional form, is autocrine. By extension, cells  
35 which produce a peptide which is being screened for the ability to activate a receptor (e.g., by activating a G protein-coupled receptor) and also express the receptor are called "autocrine cells". In some instances, such cells can also be referred to as "putative autocrine cells" since some of the cells will express peptides from the library which will not activate the receptor which is expressed. In a library of such cells, in which a  
40 multitude of different peptides are produced, it is likely that one or more of the cells will be "autocrine" in the stricter sense of the term.



5 As used herein, the term "not produced in functional form" with regard to endogenous yeast proteins is intended to encompass proteins which are not produced in functional form for any number of reasons, for example, because of a mutation to the gene which encodes the protein or a deletion, e.g., a disruption, of the gene which encodes the protein. The term "not produced in functional form" is also intended to  
10 include conditional mutations (*e.g.* temperature sensitive mutation), wherein the protein is not produced in functional form under certain conditions. The term also includes proteins (*e.g.*, in a mutant yeast cell) that are not folded correctly (*i.e.*, the tertiary structure doesn't resemble that of the protein when normally expressed in functional form).

15 As used herein the term amino acid "motif" refers to an amino acid sequence of a wild type G protein-coupled receptor which can be mutated to give the receptor altered ligand binding and/or coupling properties. The amino acid sequence motif of a G protein coupled receptor can be represented by the amino acid sequence [X<sub>1</sub>X<sub>2</sub>X<sub>3</sub>X<sub>4</sub>] (where X<sub>1</sub> is Phe, Leu, Val, or Tyr; X<sub>2</sub> is Phe, Lys, or Gln; X<sub>3</sub> is Leu, Arg, Glu, Asn,  
20 Gln, Ser, Ala, or Leu; and X<sub>4</sub> is Ala, Cys, Asp, Glu, Gly, Ser, Thr, or Tyr). Preferably the amino acid motif lies near the carboxy terminal end of a G protein coupled receptor, e.g., near the carboxy terminal end of the seventh transmembrane domain.

## II. General Overview of Assay

25 As set out above, the present invention relates to yeast cell compositions and methods for identifying effectors of modified G protein coupled receptor protein or receptor protein complexes. The instant assays are characterized by the use of a mixture of recombinant cells to sample test compounds for receptor agonists or antagonists. As described in greater detail below, the reagent cells express a heterologous modified form  
30 of a G protein coupled receptor functionally integrated into the cell and capable of transducing a detectable signal in the yeast cell. Compounds which either agonize or antagonize the receptor function can be selected and then identified based on biochemical signals produced by the receptor, or any more distal result of receptor-mediated stimulation, for example increases in endogenous mRNA expression, etc., or,  
35 in some embodiments, by the use of reporter genes responsive to such signals. In certain embodiments, the library of compounds to be tested is a library of peptides which is expressed by the yeast cells and causes stimulation in an autocrine fashion.

The ability of compounds to modulate the signal transduction activity of the target receptor can be scored for by detecting up or down-regulation of the detection  
40 signal. For example, GTPase activity, phospholipid hydrolysis, or protein phosphorylation stimulated by the receptor can be measured directly. Alternatively, the

5 use of a reporter gene can provide a readout. In any event, a statistically significant change in the detection signal can be used to facilitate isolation of compounds of interest.

10 In certain embodiments, the cells for use in the instant assays express a modified form of a heterologous G protein-coupled receptor, e.g., an IL8 or a galanin receptor, and an endogenous G protein subunit which couples to that receptor. In one embodiment, the cells are yeast cells. Preferably, the G protein coupled receptors of the present invention have been modified such that coupling of the receptor to an endogenous cell signaling pathway, e.g, the pheromone system pathway of a yeast cell, is enhanced. For example, in preferred embodiments, the yeast cells express a  
15 heterologous G protein coupled receptor which is mutated such that it displays altered ligand binding and coupling properties from the wild type receptor, thus facilitating enhanced response of that receptor to a particular ligand or enhanced coupling of the receptor in response to a ligand when expressed in the host cell. In a preferred embodiment, the host cells express a modified form of a heterologous IL-8 receptor or a modified form of a heterologous galanin receptor and a heterologous or chimeric G  
20 protein subunit. In particularly preferred embodiments, e.g., when the host cell is a yeast cell, the heterologous G protein receptor and the heterologous G protein subunit are of the same origin, e.g., mammalian. In yet another preferred embodiment, the host cells express a mutated heterologous G protein subunit.

25 In still another preferred embodiment, the host cells, e.g., yeast cells express a chimeric G protein subunit. In particularly preferred embodiments the heterologous G protein coupled receptor and the heterologous segment of the chimeric G protein subunit are derived from the same source. For example, for a mammalian G protein coupled receptor, the chimeric G protein subunit preferably comprises a portion of a yeast G  
30 protein and a portion of a mammalian G protein subunit (e.g., human or rat). In more preferred embodiments, the second amino acid sequence in the G protein subunit chimera is derived from a mammalian G protein subunit. In particularly preferred embodiments, the second amino acid sequence is derived from a human G protein subunit sequence.

35 It will further be understood that the above embodiments are not mutually exclusive. For example, in certain preferred embodiments, a yeast cell expressing a heterologous G protein coupled receptor may express a first mutated or chimeric G protein subunit and a second, different mutated or chimeric G protein subunit to enhance coupling to the heterologous receptor.

40 In certain embodiments the yeast cells also express an indicator gene that produces a detectable signal upon functional coupling of the heterologous G protein

5 coupled receptor to the G protein. In certain embodiments the indicator gene is a reporter gene construct which including a reporter gene in operative linkage with one or more transcriptional regulatory elements responsive to the signal transduction activity of the target receptor, with the level of expression of the reporter gene providing the receptor-dependent detection signal. The amount of transcription from the reporter gene  
10 may be measured using any method known to those of skill in the art. For example, specific mRNA expression may be detected using Northern blots or specific protein product may be identified by a characteristic stain or an intrinsic activity. In preferred embodiments, the gene product of the reporter is detected by an intrinsic activity associated with that product. For instance, the reporter gene may encode a gene product  
15 that, by enzymatic activity, gives rise to a detection signal based on color, fluorescence; or luminescence.

The amount of activation of the indicator gene, e.g., expression of a reporter gene, is then compared to the amount of expression in either the same cell in the absence of the test compound or it may be compared with the amount of transcription in a  
20 substantially identical cell that lacks the specific receptors or that expresses a different receptor (e.g., a different GPCR that couples to the same G protein subunit as the test receptor). A control cell may be derived from the same cells from which the recombinant cell was prepared but which had not been modified by introduction of heterologous DNA, e.g., the DNA encoding a test polypeptide. Alternatively, it may be  
25 a cell in which the specific receptors are removed. Any difference, e.g., a statistically significant difference, in the amount of transcription indicates that the test compound has in some manner altered the activity of the specific receptor.

In other preferred embodiments, the reporter gene provides a selection method such that cells in which the compound is an effector for the receptor have a growth  
30 advantage. For example the reporter could enhance cell viability, relieve the nutritional requirement of a cell, and/or provide resistance to a drug.

By using any one of these readouts, compounds which modulate signaling via the heterologous receptor can be selected. If the compound does not appear to modulate signaling via the receptor protein, the assay may be repeated and modified by the  
35 introduction of a step in which the recombinant cell is first contacted with a known activator of the target receptor to induce signal transduction from the receptor, and the compound is assayed for its ability to inhibit the activity of the receptor, e.g., to identify receptor antagonists. In yet other embodiments, compounds can be screened for members which potentiate the response to a known activator of the receptor.

40

### III. Heterologous G Protein Coupled Receptors

5 Any wild type G protein-coupled receptor comprising an amino acid motif [X<sub>1</sub>X<sub>2</sub>X<sub>3</sub>X<sub>4</sub>] (where X<sub>1</sub> is Phe, Leu, Val, or Tyr; X<sub>2</sub> is Phe, Lys, or Gln; X<sub>3</sub> is Leu, Arg, Glu, Asn, Gln, Ser, Ala, or Leu; and X<sub>4</sub> is Ala, Cys, Asp, Glu, Gly, Ser, Thr, or Tyr) lying near the carboxy terminal end of the receptor, e.g., near the carboxy terminal end of the seventh transmembrane domain, can be modified according to the invention  
10 to display altered ligand binding and/or coupling properties. For example, the IL8A receptor has been found to comprise the amino acid sequence LGFLHSCLNPIIYAFIGQN[FRNG]FLKM (SEQ ID NO: 3). The amino acid motif is seen in the [FRNG] (SEQ ID NO:4) sequence. A mutation in the Gly residue (amino acid 320 of the receptor) of this motif (to an Arg residue) has been found to result in  
15 altered ligand binding and/or coupling properties.

Other G protein coupled receptor amino acid sequences can be scanned for this same motif or can be aligned with the IL8a receptor motif. The galanin receptor has also been found to have this motif. An alignment of the IL8 and galanin receptor motifs are presented below.

20 LGFLHSCLNPIIYAFIGQN[FRNG]FLKM rIL8aR (SEQ ID NO: 3)  
LAYSNSSVNPIIYAFLSEN[FRKA]YKQV hGalR1 (SEQ ID NO: 5)  
3  
5  
1

25 Corresponding mutations can be made in the amino acid motif in other receptors to impart similar properties onto those receptors. Exemplary G protein coupled receptors having this motif are described below.

### IL8 Receptors

30 IL-8 is produced by many cell types including NK cells and T lymphocytes in response to exogenous stimuli such as polyclonal mitogens, injurious stimuli, and infectious agents, as well as proinflammatory cytokines such IL-1 and TNF. IL-8 is a chemoattractant of neutrophils, basophils, and a small proportion (10% or less) of resting CD4<sup>+</sup> and CD8<sup>+</sup> lymphocytes. IL-8 is an important mediator of inflammation. It activates neutrophil enzyme release and is also haptotactic for melanocytes and is a  
35 comitogenic stimulant of keratinocytes.

In one embodiment, the host cells of the invention express a heterologous, preferably a modified form, of an IL8 receptor functionally integrated into a signaling pathway of the host cell. In one embodiment, the host cell is a yeast cell. In another embodiment, the host cell is a mammalian cell. In preferred embodiments, the IL-8  
40 receptor is a mammalian receptor. In one aspect, the invention pertains to a recombinant

5 yeast cell comprising a modified, heterologous (e.g., human) IL8 receptor expressed in a membrane of the yeast cell such that signal transduction activity via the heterologous (e.g., human) IL8 receptor is modulated by interaction of a region of the heterologous (e.g., human) IL8 receptor with an extracellular signal and such that signal transduction is altered from that seen in wild type receptors. In one embodiment, the host cell is a  
10 yeast cell and the heterologous (e.g., human) IL8 receptor acts as a surrogate for an endogenous yeast pheromone receptor in a pheromone response pathway of the yeast cell. Modulation of the signal transduction activity of the heterologous (e.g., human) IL8 receptor by an extracellular signal in the host cell provides a detectable signal.

In one embodiment, a human G protein coupled receptor expressed by the host  
15 cells is an IL-8 receptor. In a preferred embodiment, the cell is an IL8A receptor.

### Galanin Receptors.

Galanin is a neuroendocrine peptide which is approximately 29 amino acids in  
20 length and is recognized by G protein coupled central nervous system receptors via its N-terminus.

In one embodiment, the host cells of the invention express a heterologous, and preferably a modified form of a galinin receptor, functionally integrated into a signaling pathway of the host cell. In one embodiment, the host cell is a yeast cell. In another  
25 embodiment, the host cell is a mammalian cell. In preferred embodiments, the galinin receptor is a mammalian receptor. In one aspect, the invention pertains to a recombinant yeast cell comprising a modified, heterologous (e.g., human) galinin receptor expressed in a membrane of the yeast cell such that signal transduction activity via the heterologous (e.g., human) galinin receptor is modulated by interaction of a region of the  
30 heterologous (e.g., human) galinin receptor with an extracellular signal and such that signal transduction is altered from that seen in wild type receptors. In one embodiment, the host cell is a yeast cell and the heterologous (e.g., human) galinin receptor acts as a surrogate for an endogenous yeast pheromone receptor in a pheromone response pathway of the yeast cell. Modulation of the signal transduction activity of the  
35 heterologous (e.g., human) galinin receptor by an extracellular signal in the host cell provides a detectable signal.

In one embodiment, a human G protein coupled receptor expressed by the host cells is an galinin receptor.

To facilitate expression of a heterologous (e.g., human) G protein receptor in a  
40 host cell, e.g., a yeast cell, the coding sequence for the receptor can be operatively linked to a heterologous (e.g., yeast) leader sequence. Examples of heterologous leader sequences that can be used include the leader sequence of the yeast Ste2 receptor and the

5 invertase leader sequence. Moreover, the leader sequence of yeast  $\alpha$ -factor is often used to express heterologous receptors in yeast cells. Use of heterologous leader sequences to promote the expression of the G protein coupled receptor in yeast cells is discussed in further detail in subsections below.

10 In preferred embodiments, the yeast cell expresses a mammalian, chimeric, and/or mutant G protein subunit, in addition to the heterologous (e.g., human) G protein coupled receptor. This mammalian, chimeric, and/or mutant G protein subunit can be, for example, a mammalian, chimeric, and/or mutant  $G\alpha$  subunit. Alternatively or additionally, the yeast cell can express a mammalian, chimeric, and/or mutant  $G\gamma$  subunit. The structure and use of these mammalian and chimeric G protein subunits are  
15 described in further detail in subsections below.

When the yeast cell expresses an heterologous G protein coupled receptor (e.g., a modified G protein coupled receptor), the yeast cell preferably also expresses a chimeric G protein, e.g., a  $G\alpha$  subunit that is a GPA41- $G_{\alpha i2}$  subunit. Other preferred chimeric  $G\alpha$  subunits for use with heterologous G protein coupled receptors include a GPA41-  
20  $G_{\alpha i1}$  and GPA41- $G_{\alpha i3}$  subunit. In one embodiment, the yeast cell expresses a chimeric STE18- $G\gamma$  subunit. In one embodiment, the yeast cell expresses both a chimeric  $G\alpha$  subunit and a chimeric STE18- $G\gamma$  subunit.

In one embodiment, the yeast cell preferably also expresses a mutant G protein subunit e.g., a mammalian  $G_{\alpha s}$  having a E10K (i.e., Glu to Lys substitution at position  
25 10) mutation. Other preferred chimeric  $G\alpha$  subunits for use with heterologous G protein coupled receptor include a mammalian  $G_{\alpha s}$ D229S subunit or a mammalian  $G_{\alpha s}$ E10K + D229S subunit. In other embodiments the yeast cell expresses endogenous GPA1.

In preferred embodiments, a yeast cell of the invention that expresses an G protein coupled receptor is a mutant cell having a pheromone response pathway that is  
30 desensitized at a slower rate than a wild type strain under the same conditions of continuous stimulation of the pheromone response pathway. In one embodiment, an endogenous *FAR1* gene is not functionally expressed in the yeast cell. In another embodiment, an endogenous *SST2* gene is not functionally expressed in the yeast cell. In still other embodiments, the yeast cell has a *ste14* mutation or has a *ste2* or *ste3*  
35 mutation. In yet other embodiments, an endogenous pheromone gene is not functionally expressed in the yeast cell. Various preferred mutations that can be present in the yeast cells used as host cells for expression of a heterologous receptor are described in further detail in subsections below.

To facilitate detection of signaling through the heterologous G protein coupled  
40 receptor expressed by a yeast cell of the invention, the yeast cell preferably further comprises a selectable marker that is activated by a pheromone response pathway of the yeast cell, thereby providing the detectable signal. The marker gene can be can

5 comprise, for example, a pheromone-responsive promoter operably linked to a selectable gene or a detectable gene. An example of a suitable pheromone-responsive promoter is the *FUS1* promoter. An example of suitable marker genes are the HIS 3 gene and the LacZ gene. Generation of detectable signals, and use of markers for detection thereof, are described in further detail in subsections below.

10 In another embodiment of the invention, in addition to expression of a heterologous G protein coupled receptor the yeast cells also express a heterologous polypeptide. The heterologous polypeptide is transported to a location allowing interaction with a region of the heterologous (e.g., human) receptor expressed in the cell membrane of the yeast cell. Such peptides may act at sites other than extracellular  
15 binding sites. Modulation of the signal transduction activity of the heterologous (e.g., human) receptor by the heterologous polypeptide provides a detectable signal. Thus, in this embodiment, the heterologous polypeptide and the heterologous G protein coupled receptor form an "autocrine" system in which the yeast cell expresses a heterologous polypeptide (e.g., a test polypeptide to be evaluated for receptor agonist or antagonist  
20 activity) that may stimulate the receptor expressed by that yeast cell. In certain embodiments, the heterologous polypeptide includes a signal sequence that facilitates transport of the polypeptide to a location allowing interaction with the extracellular region of the receptor. For example, the signal sequence corresponds to a leader peptide of the *Saccharomyces cerevisiae*  $\alpha$  factor or a-factor. Development and use of this  
25 "autocrine" system is described in further detail below.

Preferred yeast host cells of the invention belong to the species *Saccharomyces cerevisiae*. Examples of other suitable yeast host cells are described further in subsection IV below.

The yeast cells expressing a heterologous (e.g., human) receptor can be used in  
30 screening assays to identify modulators of the receptor. Accordingly, in one embodiment, the invention provides a method of identifying compounds which modulate a heterologous (e.g., human) receptor (e.g., an IL8 or galanin receptor) expressed by a yeast cell, comprising the steps of:

- (i) contacting a mixture of yeast cells of the invention that express a  
35 heterologous (e.g., human) receptor with a test compound;
- (ii) allowing cells within the mixture to generate a detectable signal; and
- (iii) identifying the test compound as a modulator of the heterologous (e.g., human) receptor expressed by the yeast cell.

40 Ways of monitoring changes in a detectable signal in the yeast host cells (e.g., using a reporter gene ) are described in detail in other sections of the application. Moreover, types of compounds (e.g., various libraries of compounds) that can be screened using the assay are described in detail in other sections of the application.

- 5 Examples of screening assays using yeast cells expressing a human G protein coupled receptor are described in detail in the appended Examples.

Standard techniques for preparing recombinant DNA constructs and for manipulating yeast cells and genomes (*e.g.*, by transfection and homologous recombination) can be used to create the receptor-expressing host cells for use in the invention. Exemplary constructs and techniques for making the host cells of the invention are described in further detail in the Examples.

#### IV. Host Cells

Suitable host cells for generating the subject assay include prokaryotes, yeast, or higher eukaryotic cells, including plant and animal cells, especially mammalian cells. Prokaryotes include gram negative or gram positive organisms. Examples of suitable mammalian host cell lines include the COS-7 line of monkey kidney cells (ATCC CRL 1651) (Gluzman (1981) Cell 23:175) CV-1 cells (ATCC CCL 70), L cells, C127, 3T3, Chinese hamster ovary (CHO), HeLa, HEK-293, SWISS 3T3, and BHK cell lines.

In one embodiment, the host cells are yeast cells. The host cells of the present invention may be of any species of yeast which are cultivatable and in which an exogenous receptor can be made to engage the appropriate signal transduction machinery of the host cell. Exemplary species include *Kluyverei lactis*, *Schizosaccharomyces pombe*, and *Ustilago maydis*, with *Saccharomyces cerevisiae* being preferred. Other yeast which can be used in practicing the present invention are *Neurospora crassa*, *Aspergillus niger*, *Aspergillus nidulans*, *Pichia pastoris*, *Candida tropicalis*, and *Hansenula polymorpha*. The term "yeast", as used herein, includes not only yeast in a strictly taxonomic sense, *i.e.*, unicellular organisms, but also yeast-like multicellular fungi or filamentous fungi.

#### V. Expression Systems

In general, it will be desirable that an expression vector be capable of replication in the host cell. Heterologous DNA may be integrated into the host genome, and thereafter is replicated as a part of the chromosomal DNA, or it may be DNA which replicates autonomously, as in the case of a plasmid. In the latter case, the vector will include an origin of replication which is functional in the host. In the case of an integrating vector, the vector may include sequences which facilitate integration, *e.g.*, sequences homologous to host sequences, or encoding integrases.

Appropriate cloning and expression vectors for use with bacterial, fungal, yeast, and mammalian cellular hosts are known in the art, and are described in, for example,



5 Powels et al. (*Cloning Vectors: A Laboratory Manual*, Elsevier, New York, 1985).  
Mammalian expression vectors may comprise non-transcribed elements such as an  
origin of replication, a suitable promoter and enhancer linked to the gene to be  
expressed, and other 5' or 3' flanking nontranscribed sequences, and 5' or 3'  
nontranslated sequences, such as necessary ribosome binding sites, a poly-adenylation  
10 site, splice donor and acceptor sites, and transcriptional termination sequences.

A number of vectors exist for the expression of recombinant proteins in yeast.  
For instance, YEP24, YIP5, YEP51, YEP52, pYES2, and YRP17 are cloning and  
expression vehicles useful in the introduction of genetic constructs into *S. cerevisiae*  
(see, for example, Broach et al. (1983) in *Experimental Manipulation of Gene*  
15 *Expression*, ed. M. Inouye Academic Press, p. 83, incorporated by reference herein).  
These vectors can replicate in *E. coli* due the presence of the pBR322 ori, and in *S.*  
*cerevisiae* due to the replication determinant of the yeast 2 micron plasmid. In addition,  
drug resistance markers such as ampicillin can be used. Suitable promoters for function  
in yeast include the promoters for metallothionein, 3-phosphoglycerate kinase  
20 (Hitzeman et al., *J. Biol. Chem.* 255, 2073 (1980) or other glycolytic enzymes (Hess et  
al., *J. Adv. Enzyme Req.* 7, 149 (1968); and Holland et al. *Biochemistry* 17, 4900  
(1978)), such as enolase, glyceraldehyde-3-phosphate dehydrogenase, hexokinase,  
pyruvate decarboxylase, phospho-fructokinase, glucose-6-phosphate isomerase, 3-  
phosphoglycerate mutase, pyruvate kinase, triosephosphate isomerase, phospho-glucose  
25 isomerase, and glucokinase. Suitable vectors and promoters for use in yeast expression  
are further described in R. Hitzeman et al., EPO Publn. No. 73,657. Other promoters,  
which have the additional advantage of transcription controlled by growth conditions,  
are the promoter regions for alcohol dehydrogenase 2, isocytochrome C, acid  
phosphatase, degradative enzymes associated with nitrogen metabolism, and the afore-  
30 mentioned metallothionein and glyceraldehyde-3-phosphate dehydrogenase, as well as  
enzymes responsible for maltose and galactose utilization. Finally, promoters that are  
active in only one of the two haploid mating types may be appropriate in certain  
circumstances. Among these haploid-specific promoters, the pheromone promoters  
MF $\alpha$ 1 and MF $\alpha$ 1 are of particular interest.

35 In certain embodiments, it will be desirable to control the copy number of the  
heterologous gene which is expressed. For example, in certain embodiments, expression  
of a gene (e.g., a receptor gene or a gene which results in activation of the pheromone  
response pathway) will be from a low copy number in order to better detect ligand  
induced signaling. Exemplary low copy number plasmids suitable for use in yeast cells  
40 are known in the art and include, e.g., ARS vectors or centromeric sequences (CEN)  
(See e.g., Romanos et al. 1992. *Yeast* 8:423). In other embodiments, however, the use  
of high copy number plasmids will be desirable. Exemplary high copy number plasmids

5 are also known in the art and include *E. coli*-yeast shuttle vectors based on 2 $\mu$ . In yet other embodiments it may be desirable to express heterologous DNA in a yeast cell using integrating vectors, such as YIp vectors. The use of high DNA concentrations of integrating vectors can result in tandem multicopy inserts due to repeated recombination events. Alternatively, heterologous DNA can be integrated into reiterated chromosomal  
10 DNA to generate stable multi-copy integrants (Kingsman et al. 1985. *Biotechnol. Genet. Eng. Revs.* 3:377; Lopes et al. 1989. *Gene* 79:199)

#### VI. Expression of Heterologous Receptors

For expression in yeast, a heterologous G protein coupled receptor gene can be  
15 operably linked to a promoter functional in the cell to be engineered and to a signal sequence that also functions in the cell. For example, suitable promoters include Ste2, Ste3 and gal10. Optionally, the codons of the gene can be optimized for expression in yeast. See Hoekema et al., (1987) *Mol. Cell. Biol.*, 7:2914-24; Sharp, et al., (1986) 14:5125-43.

20 In some instances a foreign receptor which is expressed in yeast will functionally integrate into the yeast membrane, and there interact with the endogenous yeast G protein. In other instances, either the receptor may be modified or a compatible G protein or a chimeric (i.e., part yeast/part mammalian), or a mutant mammalian G protein subunit which can properly interact with the exogenous receptor G protein may  
25 be provided. The homology of seven transmembrane domain receptors (STRs) is discussed in Dohlman et al., *Ann. Rev. Biochem.*, (1991) 60:653-88. When STRs are compared, a distinct spatial pattern of homology is discernible. The transmembrane domains are often the most similar, whereas the N- and C-terminal regions, and the cytoplasmic loop connecting transmembrane segments V and VI are more divergent.  
30 The functional significance of different STR regions has been studied by introducing point mutations (both substitutions and deletions) and by constructing chimeras of different but related STRs. Synthetic peptides corresponding to individual segments have also been tested for activity. Affinity labeling has been used to identify ligand binding sites. Such information can be useful in creating mutations in GPCRs to  
35 enhance functionality.

If a naturally occurring exogenous GPCR cannot be made functional in yeast, it may be mutated for this purpose. For example, a comparison can be made of the amino acid sequences of the exogenous receptor and of the yeast receptors, and regions of high and low homology identified. Trial mutations can be made to distinguish regions  
40 involved in ligand or G protein binding, from those necessary for functional integration in the membrane. The exogenous receptor can then be mutated in the latter region to

5 more closely resemble the yeast receptor, until functional integration was achieved. If  
this were insufficient to achieve functionality, mutations could next be made in the  
regions involved in G protein binding. Alternatively, the naturally occurring exogenous  
GPCR can be mutated to more closely resemble another mammalian receptor that is  
known to functionally integrate in yeast cells or random mutagenesis can be performed,  
10 followed by selection of mutants that can functionally integrate in the yeast cells.  
Another possible approach for achieving functional integration of the receptor is to make  
a chimeric receptor (mammalian/yeast)(see *e.g.*, U.S. Patent No. 5,576,210 issued to  
Sledziewski et al).

Preferably, the yeast genome is modified so that it is unable to produce the yeast  
15 receptors or G proteins which are homologous to the exogenous receptors or exogenous  
G protein subunits in functional form in order to facilitate assay interpretation. For  
example, the endogenous G protein subunit(s) is mutated generating, for instance, a  
temperature sensitive mutant.

In other embodiments, a secretory signal of a yeast protein can be used to direct  
20 transport of the heterologous G protein coupled receptors to the plasma membrane.  
Previous work has demonstrated the secretory expression of foreign proteins in yeast  
cells using the signal sequence of a yeast secreted protein such as invertase or acid  
phosphatase, encoded by the *SUC2* and *PHO5* genes, respectively (Schraber, M.D. et al.  
(1986) *Methods Enzymol.* 119:416; Moir, D.T. et al. (1991) *Methods Enzymol.* 194:491-  
25 507). The vast majority of the secreted proteins possess a hydrophobic N-terminal  
signal sequence which targets them to the endoplasmic reticulum. A leader sequence of  
the  $\alpha$ -factor precursor encoded by the *MF $\alpha$ 1* gene was shown to promote the most  
efficient secretion of various heterologous proteins. In addition to a signal sequence, the  
 $\alpha$ -factor leader includes a hydrophilic pro-region which is believed to facilitate protein  
30 transport at the later stages of the secretory pathway.

Both secreted and membrane proteins including G protein-coupled receptors are  
delivered to the cell surface through the same secretory pathway. Some receptors, for  
example, metabotropic glutamate receptors and vasoactive intestinal peptide receptors,  
also possess the N-terminal signal sequence, whereas some do not. In the latter case, a  
35 first transmembrane domain is believed to interact with the ER translocation machinery.  
The use of yeast secretory signals, in particular, the  $\alpha$ -factor leader, may be desirable to  
provide the more efficient integration of the receptors into the membrane of the  
endoplasmic reticulum and transport to the plasma membrane. In fact, the cell surface  
expression of the rat M5 receptor directed by the  $\alpha$ -factor leader has been documented  
40 (Huang et al. (1992) *Biochem. Biophys. Res. Commun.* 181:1180).

5 VII. G protein subunits and complexes

In certain instances it will be desirable to modify naturally occurring forms of yeast or mammalian G-protein subunits. For instance, where a heterologous GPCR does not adequately couple to the endogenous yeast G protein subunit, such a subunit, e.g., GPA1 may be modified to improve coupling. Such modifications can be made by  
10 mutation, e.g., directed mutation or random mutation, using methods known in the art and described in more detail below.

Alternatively, a heterologous subunit can be expressed. The specificity of coupling of a receptor to a heterotrimeric G-protein is largely determined by the  $\alpha$  subunit of the G-protein. Thus, in preferred embodiments, a heterologous  $G\alpha$  subunit is  
15 expressed in the yeast cell. The predominant role of the yeast  $G\alpha$ , GPA1, is to bind to and sequester the effector-signaling  $\beta\gamma$  component of the heterotrimer. Thus, in order to achieve functional integration into a yeast pheromone signaling pathway, a heterologous  $G\alpha$  subunit must bind to yeast  $\beta\gamma$  in the quiescent state, and release it upon receptor activation.

If functional integration is not achieved, or is not optimal, the heterologous subunit can be mutated. For example, in general, mammalian  $G\alpha$  subunits couple poorly to the  $\beta\gamma$  subunits of yeast cells. In yeast which lack their own endogenous  $G\alpha$  subunit, this failure to couple results in the constitutive activation of the pheromone pathway due to the effector activity of the unbound yeast  $\beta\gamma$ . Accordingly, if a naturally  
20 occurring heterologous G protein subunit does not enhance coupling, modifications can be made. Such modifications may take the form of mutations which are designed to increase the resemblance of the G protein subunit to the yeast G protein subunit while decreasing its resemblance to the heterologous receptor-associated G protein subunit.

For example, a residue may be changed so as to become identical to the  
30 corresponding yeast G protein residue, or to belong to the same exchange group of that residue. After modification, the modified G protein subunit might or might not be "substantially homologous" to the heterologous and/or the yeast G protein subunit.

In the case of  $G\alpha$ , modifications are preferably concentrated in regions of the  $G\alpha$  which are likely to be involved in  $G\beta\gamma$  binding.

In other embodiments, modifications will take the form of replacing one or more  
35 amino acids of the receptor-associated G protein subunit with the corresponding yeast G protein subunit amino acids, thereby forming a chimeric G protein subunit. In preferred embodiments, three or more consecutive amino acids are replaced. In other embodiments, point mutations may be sufficient.

5 Chimeric G protein subunits of the invention enhance coupling of the heterologous receptor to the endogenous yeast signaling pathway. For example, a chimeric  $G\alpha$  subunit will interact with the heterologous receptor and the yeast  $G\beta\gamma$  complex, thereby permitting signal transduction.

10 A yeast cell of the present invention can express one or more of the indicated G protein structures. For example, a yeast cell can express a chimeric or mutant  $G\alpha$  subunit, and an endogenous yeast  $G\beta\gamma$ , a mammalian  $G\beta\gamma$ , a mutated mammalian  $G\beta\gamma$ , or a chimeric  $G\beta\gamma$ .

15 In preferred embodiments, both the receptor and the heterologous subunit are derived from the same source, e.g., are mammalian. In particularly preferred embodiment, both are human in origin.

In another preferred embodiment, a yeast cell that expresses a heterologous or chimeric G protein subunit has been modified such that the endogenous, homologous G protein subunit gene is disrupted.

20 In certain embodiments, yeast strains lacking pheromone receptors and having no heterologous receptor capable of coupling to the pheromone response pathway may be used to assess the affinity of an endogenous yeast G protein subunit, a mammalian G protein subunit, a mutated G protein subunit, or chimeric G protein subunit for other yeast subunits. For example, the affinity of GPA1p, chimeric GPA- $G\alpha$ s, or other  $G\alpha$  subunit for yeast  $\beta\gamma$  or other chimeric  $\beta\gamma$  subunit can be assessed. Such strains depend  
25 on free  $\beta\gamma$  for signaling through the pheromone response pathway leading to growth arrest. Mutant  $G\alpha$  subunits may be tested in such a system, those which bind  $\beta\gamma$  more effectively will sequester  $\beta\gamma$  and reduce or block signaling. Preferably, such chimeras and GPA1 subunits can be assessed in a *gpa1*<sup>-</sup> background to avoid competition with GPA1 for  $\beta\gamma$ . For example,  $G\alpha$ s chimeric mutants (see below) carrying D229S, E10K, N254D, or S286P were found to sequester  $\beta\gamma$  more effectively than the mutant with wild  
30 type sequences. Also, double mutants were even more effective than either single mutant. Similarly, overexpression of  $G\alpha$ s by driving transcription from the highly efficient PGK promoter resulted in dampening of the receptor coupling which may be offset by introduction of the double mutant  $G\alpha$ s (D229S,E10K).

35 Guidance for making mutations in G protein subunits and in the construction of chimeric G protein subunits is provided below.

#### *Site-Directed Mutagenesis versus Random Mutagenesis*

40 There are numerous art recognized ways to solve the structure-function problems of the sort presented by attempts to define the determinants involved in mediating the

5 association of the subunits that comprise the G protein heterotrimer. For example, in one approach, discussed above with respect to hybrid constructs, specific mutations or alterations are introduced into a molecule based upon the available experimental evidence. In a second approach, random mutagenesis techniques, coupled with selection or screening systems, are used to introduce large numbers of mutations into a molecule, and that collection of randomly mutated molecules is then subjected to a selection for the desired phenotype or a screen in which the desired phenotype can be observed against a background of undesirable phenotypes.

15 With random mutagenesis one can mutagenize an entire molecule or one can proceed by cassette mutagenesis. In the former instance, the entire coding region of a molecule is mutagenized by one of several methods (chemical, PCR, doped oligonucleotide synthesis) and that collection of randomly mutated molecules is subjected to selection or screening procedures. Random mutagenesis can be applied in this way in cases where the molecule being studied is relatively small and there are powerful and stringent selections or screens available to discriminate between the different classes of mutant phenotypes that will inevitably arise. In the second approach, discrete regions of a protein, corresponding either to defined structural (i.e.  $\alpha$ -helices,  $\beta$ -sheets, turns, surface loops) or functional determinants (e.g., catalytic clefts, binding determinants, transmembrane segments) are subjected to saturating or semi-random mutagenesis and these mutagenized cassettes are re-introduced into the context of the otherwise wild type allele.

Cassette mutagenesis is most useful when there is experimental evidence available to suggest a particular function for a region of a molecule and there is a powerful selection and/or screening approach available to discriminate between interesting and uninteresting mutants. Cassette mutagenesis is also useful when the parent molecule is comparatively large and the desire is to map the functional domains of a molecule by mutagenizing the molecule in a step-wise fashion, i.e. mutating one linear cassette of residues at a time and then assaying for function.

The present invention provides for applying random mutagenesis in order to further delineate the determinants involved in  $G\alpha$ - $G\beta\gamma$  or subunit-receptor association. Random mutagenesis may be accomplished by many means, including:

1. PCR mutagenesis, in which the error prone Taq polymerase is exploited to generate mutant alleles of G protein subunits, which are assayed directly in yeast for an ability to couple.
2. Chemical mutagenesis, in which expression cassettes encoding G protein subunits are exposed to mutagens and the protein products of the mutant sequences are assayed directly in yeast for an ability to couple.

5           3. Doped synthesis of oligonucleotides encoding portions of the G protein subunit gene.

          4. *In vivo* mutagenesis, in which random mutations are introduced into the coding region of G protein subunits by passage through a mutator strain of *E. coli*, XL1-Red (mutD5 mutS mutT) (Stratagene, Menasa, WI).

10           In certain embodiments, for example, the random mutagenesis may be focused on regions suspected to be involved in  $G\alpha$ - $G\beta\gamma$  association. Random mutagenesis approaches are feasible for two reasons. First, in yeast one has the ability to construct stringent screens and facile selections (growth vs. death, transcription vs. lack of transcription) that are not readily available in mammalian systems. Second, when using  
15           yeast it is possible to screen efficiently through thousands of transformants rapidly. For example, this relatively small region of  $G\alpha$  subunits represents a reasonable target for cassette mutagenesis. Another region that may be amenable to cassette mutagenesis is that defining the surface of the switch region of  $G\alpha$  subunits that is solvent-exposed in the crystal structures of  $G\alpha_i$  and transducin. From the data described below, this surface  
20           may contain residues that are in direct contact with yeast  $G\beta\gamma$  subunits, and may therefore be a reasonable target for mutagenesis.

#### *A. Modification of $G\alpha$*

          Some aspects of  $G\alpha$  structure are relevant to the design of modified  $G\alpha$  subunits.

25           Alignments of  $G\alpha$  and GPA1 can be made to determine sequence similarity. For alignments of the entire coding regions of GPA1 with  $G\alpha_s$ ,  $G\alpha_i$ , and  $G\alpha_o$ ,  $G\alpha_q$  and  $G\alpha_z$ , see Dietzel and Kurjan (1987, *Cell* 50:573) and Lambright, et al. (1994, *Nature* 369:621-628). Additional sequence information is provided by Mattera, et al. (1986, *FEBS Lett* 206:36-41), Bray, et al. (1986, *Proc. Natl. Acad. Sci USA* 83:8893-8897) and  
30           Bray, et al. (1987, *Proc Natl. Acad Sci USA* 84:5115-5119). An alignment of GPA1 and four other  $G\alpha$  proteins is provided by Stone and Reed (1990. *Mol. Cell Biol.* 10:4439).

          The gene encoding a G protein homolog of *S. cerevisiae* was cloned independently by Dietzel and Kurjan (*supra*) (who referred to the gene as SCG1) and by Nakafuku, et al. (1987 *Proc Natl Acad Sci* 84:2140-2144) (who called the gene GPA1).  
35           Sequence analysis revealed a high degree of homology between the protein encoded by this gene and mammalian  $G\alpha$ . GPA1 encodes a protein of 472 amino acids, as compared with approximately 340-350 amino acids for most mammalian  $G\alpha$  subunits in four described families,  $G\alpha_s$ ,  $G\alpha_i$ ,  $G\alpha_q$  and  $G\alpha_{12/13}$ . Nevertheless, GPA1 shares overall sequence and structural homology with all  $G\alpha$  proteins identified to date. The highest  
40           overall homology in GPA1 is to the  $G\alpha_i$  family (48% identity, or 65% with conservative

5 substitutions) and the lowest is to G $\alpha$ s (33% identity, or 51% with conservative substitutions) (Nakafuku, et al., *supra*).

The regions of high sequence homology among G $\alpha$  subunits are dispersed throughout their primary sequences, with the regions sharing the highest degree of homology mapping to sequence that comprises the guanine nucleotide binding/GTPase  
10 domain. This domain is structurally similar to the  $\alpha\beta$  fold of ras proteins and the protein synthesis elongation factor EF-Tu. This highly conserved guanine nucleotide-binding domain consists of a six-stranded  $\beta$  sheet surrounded by a set of five  $\alpha$ -helices. It is within these  $\beta$  sheets and  $\alpha$  helices that the highest degree of conservation is observed among all G $\alpha$  proteins, including GPA1. The least sequence and structural homology is  
15 found in the intervening loops between the  $\beta$  sheets and  $\alpha$  helices that define the core GTPase domain. There are a total of four "intervening loops" or "inserts" present in all G $\alpha$  subunits. In the crystal structures reported to date for the GDP- and GTP $\gamma$ S-liganded forms of bovine rod transducin (Noel, et al. (1993) *Nature* 366:654-663); (Lambright, et al. (1994) *Nature* 369:621-628), the loop residues are found to be outside the core  
20 GTPase structure. Functional roles for these loop structures have been established in only a few instances. A direct role in coupling to phosphodiesterase- $\gamma$  has been demonstrated for residues within inserts 3 and 4 of G $\alpha$ t (Rarick, et al. (1992) *Science* 256:1031-1033); (Artemyev, et al. (1992) *J. Biol. Chem.* 267:25067-25072), while a "GAP-like" activity has been ascribed to the largely  $\alpha$ -helical insert 1 domain of G $\alpha$ S  
25 (Markby, et al. (1993) *Science* 262:1805-1901).

While the amino- and carboxy-termini of G $\alpha$  subunits do not share striking homology either at the primary, secondary, or tertiary levels, there are several generalizations that can be made about them. First, the amino termini of G $\alpha$  subunits have been implicated in the association of G $\alpha$  with G $\beta\gamma$  complexes and in membrane  
30 association via N-terminal myristoylation. In addition, the carboxy-termini have been implicated in the association of G $\alpha\beta\gamma$  heterotrimeric complexes with G protein-coupled receptors (Sullivan, et al. (1987) *Nature* 330:758-760); West, et al. (1985) *J. Biol. Chem.* 260:14428-14430); (Conklin, et al. (1993) *Nature* 363:274-276); (Kallal and Kurjan. 1997. *Mol. Cell. Biol.* 17:2897). Data in support of these generalizations about the  
35 function of the N-terminus derive from several sources, including both biochemical and genetic studies.

In the GPA<sub>41</sub>G $\alpha$  hybrids, the amino terminal 41 residues are derived from GPA1. All residues following position 41 are contributed by the human G $\alpha$  subunits, including the consensus nucleotide binding motif. For alignments of the entire coding regions of  
40 GPA1 with G $\alpha$ s, G $\alpha$ i, and G $\alpha$ O, G $\alpha$ q and G $\alpha$ z, see Dietzel and Kurjan (1987, *Cell* 50:573) and Lambright, et al. (1994, *Nature* 369:621-628). Additional sequence



5 information is provided by Mattera, et al. (1986, *FEBS Lett* 206:36-41), Bray, et al. (1986, *Proc. Natl. Acad. Sci USA* 83:8893-8897) and Bray, et al. (1987, *Proc Natl. Acad Sci USA* 84:5115-5119).

There is little if any sequence homology shared among the amino termini of  $G\alpha$  subunits. The amino terminal domains of  $G\alpha$  subunits that precede the first  $\beta$ -sheet vary  
10 in length from 41 amino acids (GPA1) to 31 amino acids ( $G\alpha_t$ ). Most  $G\alpha$  subunits share the consensus sequence for the addition of myristic acid at their amino termini, although not all  $G\alpha$  subunits that contain this motif have myristic acid covalently associated with the glycine at position 2 (Speigel, et al. (1991) *TIBS* 16:338-3441). The role of this post-translational modification has been inferred from studies in which the activity of mutant  
15  $G\alpha$  subunits from which the consensus sequence for myristoylation has been added or deleted has been assayed (Mumby et al. (1990) *Proc. Natl. Acad. Sci. USA* 87: 728-732; (Linder, et al. (1991) *J. Biol Chem.* 266:4654-4659); Gallego, et al. (1992) *Proc. Natl. Acad. Sci. USA* 89:9695-9699). These studies suggest two roles for N-terminal myristoylation. First, the presence of amino-terminal myristic acid has in some cases  
20 been shown to be required for association of  $G\alpha$  subunits with the membrane, and second, this modification has been demonstrated to play a role in modulating the association of  $G\alpha$  subunits with  $G\beta\gamma$  complexes. The role of myristoylation of the GPA1 gene products, at present is unknown.

In other biochemical studies aimed at examining the role of the amino-terminus  
25 of  $G\alpha$  in driving the association between  $G\alpha$  and  $G\beta\gamma$  subunits, proteolytically or genetically truncated versions of  $G\alpha$  subunits were assayed for their ability to associate with  $G\beta\gamma$  complexes, bind guanine nucleotides and/or to activate effector molecules. In all cases,  $G\alpha$  subunits with truncated amino termini were deficient in all three functions (Graf, et al. (1992) *J. Biol. Chem.* 267:24307-24314); (Journot, et al. (1990) *J. Biol.*  
30 *Chem.* 265:9009-9015); and (Neer, et al. (1988) *J. Biol. Chem* 263:8996-9000). Slepak, et al. (1993, *J. Biol. Chem.* 268:1414-1423) reported a mutational analysis of the N-terminal 56 a.a. of mammalian  $G\alpha_o$  expressed in *Escherichia coli*. Molecules with an apparent reduced ability to interact with exogenously added mammalian  $G\beta\gamma$  were identified in the mutant library. As the authors pointed out, however, the assay used to  
35 screen the mutants the extent of ADP-ribosylation of the mutant  $G\alpha$  by pertussis toxin was not a completely satisfactory probe of interactions between  $G\alpha$  and  $G\beta\gamma$ . Mutations identified as inhibiting the interaction of the subunits, using this assay, may still permit the complexing of  $G\alpha$  and  $G\beta\gamma$  while sterically hindering the ribosylation of  $G\alpha$  by toxin. Other work has revealed specific amino acid residues of GPA1 that are important  
40 in GPA1 function. For example, a E307K mutation appears to create an  $\alpha$  subunit with a broadened specificity for  $G\beta$  subunits (Whiteway et al. 1994. *Mol. Cell. Biol.* 14:3223). Interestingly, the residue in the mammalian  $G\alpha$  subunit which is equivalent

5 to the E307 position is diagnostic for a particular class of mammalian  $\alpha$  subunits. For example, the  $G_s\alpha$  subunits contain a lysine at this position, the  $G_o$  and  $G_i$   $\alpha$  subunits contain a histidine, the transducin  $\alpha$  subunits have a glutamine, the  $G_q$   $\alpha$  subunits have a proline, and the  $G_{13}$   $\alpha$  subunits have an aspartic acid at this site (Whiteway et al. *supra*).

Genetic studies examined the role of amino-terminal determinants of  $G\alpha$  in  
10 heterotrimer subunit association have been carried out in both yeast systems using GPA1-mammalian  $G\alpha$  hybrids (Kang, et al. (1990) *Mol. Cell. Biol.* 10:2582-2590) and in mammalian systems using  $G\alpha_i/G\alpha_s$  hybrids (Russell and Johnson (1993) *Mol. Pharmacol.* 44:255-263). In the former studies, gene fusions, composed of yeast GPA1 and mammalian  $G\alpha$  sequences were constructed by Kang, et al. (*supra*) and assayed for  
15 their ability to complement a *gpa1* null phenotype (i.e., constitutive activation of the pheromone response pathway) in *S. cerevisiae*. Kang, et al. demonstrated that wild type mammalian  $G\alpha_s$ ,  $G\alpha_i$  but not  $G\alpha_o$  proteins are competent to associate with yeast  $G\alpha$  and suppress the *gpa1* null phenotype, but only when overexpressed. Fusion proteins containing the amino-terminal 330 residues of GPA1 sequence linked to 160, 143, or 142  
20 residues of the mammalian  $G\alpha_s$ ,  $G\alpha_i$  and  $G\alpha_o$  carboxyl-terminal regions, respectively, also coupled to the yeast mating response pathway when overexpressed on high copy plasmids with strong inducible (CUP) or constitutive (PGK) promoters. All three of these hybrid molecules were able to complement the *gpa1* null mutation in a growth arrest assay, and were additionally able to inhibit  $\alpha$ -factor responsiveness and mating in  
25 tester strains. These last two observations argue that hybrid yeast-mammalian  $G\alpha$  subunits are capable of interacting directly with yeast  $G\beta\gamma$ , thereby disrupting the normal function of the yeast heterotrimer. Fusions containing the amino terminal domain of  $G\alpha_s$ ,  $G\alpha_i$  or  $G\alpha_o$ , however, did not complement the *gpa1* null phenotype, indicating a requirement for determinants in the amino terminal 330 amino acid residues  
30 of GPA1 for association and sequestration of yeast  $G\beta\gamma$  complexes. Taken together, these data suggest that determinants in the amino terminal region of  $G\alpha$  subunits determine not only the ability to associate with  $G\beta\gamma$  subunits in general, but also with specific  $G\beta\gamma$  subunits in a species-restricted manner.

Hybrid  $G\alpha_i/G\alpha_s$  subunits have been assayed in mammalian expression systems  
35 (Russell and Johnson (*supra*)). In these studies, a large number of chimeric  $G\alpha$  subunits were assayed for an ability to activate adenylyl cyclase, and therefore, indirectly, for an ability to interact with  $G\beta\gamma$  (i.e., coupling of  $G\alpha$  to  $G\beta\gamma$ = inactive cyclase; uncoupling of  $G\alpha$  from  $G\beta\gamma$  = active cyclase). From these studies a complex picture emerged in which determinants in the region between residues 25 and 96 of the hybrids were found to  
40 determine the state of activation of these alleles as reflected in their rates of guanine nucleotide exchange and GTP hydrolysis and the extent to which they activated adenylyl cyclase *in vivo*. These data could be interpreted to support the hypothesis that structural

5 elements in the region between the amino terminal methionine and the  $\beta$  sheet identified  
in the crystal structure of G $\alpha$ t (see Noel, et al. *supra* and Lambright, et al. *supra*) are  
involved in determining the state of activity of the heterotrimer by (1) driving  
association/dissociation between G $\alpha$  and G $\beta\gamma$  subunits; (2) driving GDP/GTP exchange.  
While there is no direct evidence provided by these studies to support the idea that  
10 residues in this region of G $\alpha$  and residues in G $\beta\gamma$  subunits contact one another, the data  
nonetheless provide a positive indication for the construction of hybrid G $\alpha$  subunits that  
retain function. There is, however, a negative indicator that derives from this work in  
that some hybrid constructs resulted in constitutive activation of the chimeric proteins  
(i.e., a loss of receptor-dependent stimulation of G $\beta\gamma$  dissociation and effector  
15 activation).

### *B. Construction of chimeric G $\alpha$ subunits.*

In preferred embodiments chimeric G $\alpha$  subunits retain as much of the sequence  
of the native mammalian proteins as possible and, in particularly preferred  
20 embodiments, the level of expression for the heterologous components should approach,  
as closely as possible, the level of their endogenous counterparts. The results described  
by King, et al. (1990, Science 250:121-123) for expression of the human  $\beta$ 2-adrenergic  
receptor and G $\alpha$ s in yeast, taken together with negative results obtained by Kang, et al.  
(*supra*) with full-length mammalian G $\alpha$  subunits other than G $\alpha$ s, led to the following  
25 preferred embodiments for the development of yeast strains in which mammalian G  
protein-coupled receptors could be linked to the pheromone response pathway.

In one embodiment, mammalian G $\alpha$  subunits are expressed using the native  
sequence of each subunit or, alternatively, as minimal gene fusions with sequences from  
the amino-terminus of GPA1 replacing the homologous residues from the mammalian  
30 G $\alpha$  subunits. In another embodiment, mammalian G $\alpha$  subunits are expressed from the  
GPA1 promoter either on low copy plasmids or after integration into the yeast genome  
as a single copy gene. In certain embodiments, endogenous G $\beta\gamma$  subunits are provided  
by the yeast STE4 and STE18 loci, while in other embodiments chimeric or  
heterologous G $\beta$  and/or G $\gamma$  subunits are also provided.

35

### *C. Rational Design of Chimeric G $\alpha$ Subunits*

Several classes of rationally designed GPA1-mammalian G $\alpha$  hybrid subunits  
have been tested for the ability to couple to yeast  $\beta\gamma$ . The first, and largest, class of  
hybrids are those that encode different lengths of the GPA1 amino terminal domain in  
40 place of the homologous regions of the mammalian G $\alpha$  subunits. This class of hybrid

5 molecules includes GPABAMH1, GPA41, GPAID, and GPALW hybrids, described below. The rationale for constructing these hybrid G $\alpha$  proteins is based on results, described above, that bear on the importance of the amino terminal residues of G $\alpha$  in mediating interaction with G $\beta\gamma$ .

10 Preferably, the yeast G $\alpha$  subunit is replaced by a chimeric G $\alpha$  subunit in which a portion, e.g., at least about 20, more preferably at least about 40, amino acids, from the amino terminus of the yeast G $\alpha$ , is fused to a sequence from a mammalian (or other exogenous) G $\alpha$ . While about 40 amino acids is the suggested starting point, shorter or longer portions may be tested to determine the minimum length required for coupling to yeast G $\beta\gamma$  and the maximum length compatible with retention of coupling to the  
15 exogenous receptor. It is presently believed that only the final 10 or 20 amino acids at the carboxy terminus of the G $\alpha$  subunit are required for interaction with the receptor.

#### *i. GPABAMH1 hybrids.*

20 Kang et al. *supra.* described hybrid G $\alpha$  subunits encoding the amino terminal 310 residues of GPAI fused to the carboxyl terminal 160, 143 and 142 residues, respectively, of G $\alpha$ S, G $\alpha$ i2, and G $\alpha$ o. In all cases examined by Kang et al., the hybrid proteins were able to complement the growth arrest phenotype of *gal* strains. Hybrids between GPAI and G $\alpha$ i3, G $\alpha$ q and G $\alpha$ 16 can be constructed, as described below, and functionally complement the growth arrest phenotype of *gal* strains.

25 GPA41 hybrids: The rationale for constructing a minimal hybrid encoding only 41 amino acids of GPAI relies upon the biochemical evidence for the role of the amino-terminus of G $\alpha$  subunits discussed above, together with the following observation. G $\beta$  and G $\gamma$  subunits are known to interact via  $\alpha$ -helical domains at their respective amino-termini (Pronin, et al. (1992) *Proc. Natl. Acad. Sci. USA* 89:6220-6224);  
30 Garritsen, et al. 1993). The suggestion that the amino termini of G $\alpha$  subunits may form an helical coil and that this helical coil may be involved in association of G $\alpha$  with G $\beta\gamma$  (Masters et al (1986) *Protein Engineering* 1:47-54); Lupas et al.(1992) *FEBS Lett.* 314:105-108) leads to the hypothesis that the three subunits of the G-protein heterotrimer interact with one another reversibly through the winding and unwinding of  
35 their amino-terminal helical regions. (For further description of the crystal structure of G proteins, and subunits thereof, see Lambright et al. (1996) *Nature* 379:311-319 and Sondek et al. (1996) *Nature* 379:369-374.) A mechanism of this type has been suggested, as well, from an analysis of leucine zipper mutants of the GCN4 transcription factor (Harbury, et al. (1993) *Science* 262:1401-1407). The rationale for constructing  
40 hybrids like those described by Kang, et al. *supra.*, that contain a majority of yeast sequence and only minimal mammalian sequence, derives from their ability to function

5 in assays of coupling between  $G\alpha$  and  $G\beta\gamma$  subunits. However, these chimeras had never been assayed for an ability to couple to both mammalian G protein-coupled receptors and yeast  $G\beta\gamma$  subunits, and hence to reconstitute a hybrid signaling pathway in yeast.

10 GPA<sub>41</sub> hybrids that have been constructed and tested include  $G\alpha_s$ ,  $G\alpha_i2$ ,  $G\alpha_i3$ ,  $G\alpha_q$ ,  $G\alpha_{o_a}$ ,  $G\alpha_{o_b}$  and  $G\alpha_{16}$ . Hybrids of  $G\alpha_s$ ,  $G\alpha_i2$ ,  $G\alpha_i3$ , and  $G\alpha_{16}$  functionally complement the growth arrest phenotype of *gpa1* strains, while GPA<sub>41</sub> hybrids of  $G\alpha_{o_a}$  and  $G\alpha_{o_b}$  do not. In addition to being tested in a growth arrest assay, these constructs have been assayed in the more sensitive transcriptional assay for activation of a *fuslp-HIS3* gene. In both of these assays, the GPA<sub>41</sub>- $G\alpha_s$  hybrid couples less well than the  
15 GPA<sub>41</sub>-i2, -i3, and -16 hybrids, while the GPA<sub>41</sub> - $\alpha_a$ , and - $\alpha_b$  hybrids do not function in either assay.

Several predictive algorithms indicate that the amino terminal domain up to the highly conserved sequence motif LLLLGAGESG (the first L in this motif is residue 43 in GPA1) forms a helical structure with amphipathic character. Assuming that a  
20 heptahelical repeat unit, the following hybrids between yeast GPA1 and mammalian  $G\alpha_s$  can be used to define the number of helical repeats in this motif necessary for hybrid function:

GPA1-7/ $G\alpha_s$ 8-394

GPA1-14/ $G\alpha_s$ 15-394

25 GPA1-21/ $G\alpha_s$ 22-394

GPA1-28/ $G\alpha_s$ 29-394

GPA1-35/ $G\alpha_s$ 36-394

GPA1-42/ $G\alpha_s$ 43-394

In these hybrids, the prediction is that the structural repeat unit in the amino  
30 terminal domain up to the tetra-leucine motif is 7, and that swapping sequences in units of 7 will in effect amount to a swap of unit turns of turns of the helical structure that comprises this domain.

A second group of "double crossover" hybrids of this class are those that are aligned on the first putative heptad repeat beginning with residue G11 in GPA1. In  
35 these hybrids, helical repeats are swapped from GPA1 into a  $G\alpha_s$  backbone one heptad repeat unit at a time.

$G\alpha_s$ 1-10/GPA11-17/ $G\alpha_s$ 18-394

$G\alpha_s$ 1-17/GPA18-24/ $G\alpha_s$ 25-394

5 GαS1-17/GPA25-31/GαS32-394

GαS1-17/GPA32-38/GαS39-394

~ The gap that is introduced between residues 9 and 10 in the GαS sequence is to preserve the alignment of the LLLLGAGE sequence motif. This class of hybrids can be complemented by cassette mutagenesis of each heptad repeat followed by screening of these collections of "heptad" libraries in standard coupling assays.

A third class of hybrids based on the prediction that the amino terminus forms a helical domain with a heptahelical repeat unit are those that effect the overall hydrophobic or hydrophilic character of the opposing sides of the predicted helical structure (See Lupas et al. *supra*). In this model, the a and d positions of the heptad repeat abcdefg are found to be conserved hydrophobic residues that define one face of the helix, while the e and g positions define the charged face of the helix. In this class of hybrids, the sequence of the GαS parent is maintained except for specific substitutions at one or more of the following critical residues to render the different helical faces of GαS more "GPA1-like"

20

K8Q

+I-10

E10G

Q12E

25

R13S

N14D

E15P

E15F

K17L

30

E21R

K28Q

K32L

V36R

This collection of single mutations could be screened for coupling efficiency to yeast Gβγ and then constructed in combinations (double and greater if necessary).

A fourth class of hybrid molecules that span this region of GPA1-Gα hybrids are those that have junctions between GPA1 and Gα subunits introduced by three primer PCR. In this approach, the two outside primers are encoded by sequences at the initiator methionine of GPA1 on the 5' side and at the tetra-leucine motif of GαS (for example) on the 3' side. A series of junctional primers spanning different junctional points can be

5 mixed with the outside primers to make a series of molecules each with different amounts of GPA1 and G $\alpha$ S sequences, respectively.

ii. *GPA<sub>ID</sub> and GPA<sub>LW</sub> hybrids.*

a  
The regions of high homology among G $\beta\gamma$  subunits that have been identified by  
10 sequence alignment are interspersed throughout the molecule. The G1 region containing the highly conserved GSGESGDST motif is followed immediately by a region of very low sequence conservation, the "il" or insert 1 region. Both sequence and length vary considerably among the il regions of the G $\alpha$  subunits. By aligning the sequences of G $\alpha$  subunits, the conserved regions bounding the il region were identified and two  
15 additional classes of GPA1-G $\alpha$  hybrids were constructed. The GPA<sub>ID</sub> hybrids encode the amino terminal 102 residues of GPA1 (up to the sequence QARKLGIQ) fused in frame to mammalian G $\alpha$  subunits, while the GPA<sub>LW</sub> hybrids encode the amino terminal 244 residues of GPA1. The reason for constructing the GPA<sub>ID</sub> and GPA<sub>LW</sub> hybrids was to test the hypothesis that the il region of GPA1 is required for mediating the  
20 interaction of GPA1 with yeast G $\beta\gamma$  subunits, for the stable expression of the hybrid molecules, or for function of the hybrid molecules. The GPA<sub>ID</sub> hybrids contain the amino terminal domain of GPA1 fused to the il domain of mammalian subunits, and therefore do not contain the GPA1 il region, while the GPA<sub>LW</sub> hybrids contain the amino terminal 244 residues of GPA1 including the entire il region (as defined by  
25 sequence alignments). Hybrids of both GPA<sub>ID</sub> and GPA<sub>LW</sub> classes were constructed for G $\alpha$ S, C- $\alpha$ i2, G $\alpha$ i3, G $\alpha$ o<sub>a</sub>, and G $\alpha$ 16; none of these hybrids complemented the gpal growth arrest phenotype.

a  
Subsequent to the construction and testing of the GPA<sub>ID</sub> and GPA<sub>LW</sub> classes of hybrids, the crystal structures of G<sub>transducin</sub> in both the GDP and GTP $\gamma$ S-liganded  
30 form, and the crystal structure of several G $\alpha$ il variants in the GTP $\gamma$ S-liganded and GDP-AlF<sub>4</sub> forms were reported (Noel et al. *supra*; Lambright et al. *supra*; and Coleman et al.(1994) *Science* 265:1405-1412). The crystal structures reveal that the il region defined by sequence alignment has a conserved structure that is comprised of six alpha helices in a rigid array, and that the junctions chosen for the construction of the GPA<sub>ID</sub> and GPA<sub>LW</sub> hybrids were not compatible with conservation of the structural features of  
35 the il region observed in the crystals. The junction chosen for the GPA<sub>ID</sub> hybrids falls in the center of the long  $\alpha$ A helix; chimerization of this helix in all likelihood destabilizes it and the protein structure in general. The same is true of the junction chosen for the GPA<sub>LW</sub> hybrids in which the crossover point between GPA1 and the  
40 mammalian G $\alpha$  subunit falls at the end of the short  $\alpha$ C helix and therefore may distort it and destabilize the protein.

The failure of the GPA<sub>ID</sub> and GPA<sub>LW</sub> hybrids is predicted to be due to disruption of critical structural elements in the il region as discussed above. Based upon new alignments and the data presented in Noel et al (*supra*), Lambright et al (*supra*), and Coleman et al (*supra*), this problem can be averted with the ras-like core domain and the il helical domain are introduced outside of known structural elements like alpha-helices.

Hybrid A GαS1-67/GPA66-299/GαS203-394

This hybrid contains the entire il insert of GPA1 interposed into the GαS sequence.

Hybrid B GPA1-41/GαS4443-67/GPA66-299/GαS203-394

This hybrid contains the amino terminal 41 residues of GPA1 in place of the 42 amino terminal residues of GαS found in Hybrid A.

### iii. Gαs Hybrids.

There is evidence that the "switch region" encoded by residues 171-237 of Gα transducin (using the numbering of (Noel et al (*supra*))) also plays a role in Gβγ coupling. First, the G226A mutation in GαS prevents the GTP-induced conformational change that occurs with exchange of GDP for GTP upon receptor activation by ligand. This residue maps to the highly conserved sequence DVGGQ, present in all Gα subunits and is involved in GTP hydrolysis. In both the Gαt and Gαi crystal structures, this sequence motif resides in the loop that connects the β3 sheet and the α2 helix in the guanine nucleotide binding core. In addition to blocking the conformational change that occurs upon GTP binding, this mutation also prevents dissociation of GTP-liganded Gαs from Gβγ. Second, crosslinking data reveals that a highly conserved cysteine residue in the α2 helix (C215 in Gαo, C210 in Gαt) can be crosslinked to the carboxy terminal region of Gβ subunits. Finally, genetic evidence (Whiteway et al. (1993) Mol Cell Biol. 14:3233-3239) identifies an important single residue in GPA1 (E307) in the β2 sheet of the core structure that may be in direct contact with βγ. A mutation in the GPA1 protein at this position suppresses the constitutive signaling phenotype of a variety of STE4 (Gβ) dominant negative mutations that are also known to be defective in Gα-Gβγ association (as assessed in two-hybrid assay in yeast as well as by more conventional genetic tests).

The GPA1 switch region suppresses coupling to yeast Gβγ (SGS), while in the context of the GPA1 amino terminus the GPA1 switch region stabilizes coupling with Gβ (Gβ<sub>2</sub> SGS). This suggests that the switch region of GPA1 stabilizes coupling with



- 5 mitigated by the observation that the GPA<sub>41</sub>-G $\alpha$ S hybrid that does not contain the GPA1 switch region is able to complement the growth arrest phenotype of gpa1 strains.

The role of the surface-exposed residues of this region may be crucial for effective coupling to yeast G $\beta\gamma$ , and can be incorporated into hybrid molecules as follows below.

10

G $\alpha$ S-GPA-Switch      G $\alpha$ S 1-202/GPA298-350/G $\alpha$ S 253-394

This hybrid encodes the entire switch region of GPA 1 in the context of G $\alpha$ S.

G $\alpha$ S-GPA- $\alpha$ 2      GQS 1-226/GPA322-332/GQS 238-394

15

This hybrid encodes the  $\alpha$ 2 helix of GPA1 in the context of G $\alpha$ S.

GPA41-G $\alpha$ S-GPA- $\alpha$ 2GPA1-41/GQS43-226/GPA322-332/GQS238-394

This hybrid encodes the 41 residue amino terminal domain of GPA1 and the  $\alpha$ 2 helix of GPA1 in the context of G $\alpha$ S.

20

In addition, hybrids that alter the surface exposed residues of the  $\beta$ 2 and  $\beta$ 3 sheets of  $\alpha$ S so that they resemble those of the GPA1 QS helix can be made. These altered  $\alpha$ 2 helical domains have the following structure. (The positions of the altered residues correspond to G $\alpha$ S.)

25

L203K

K211E

D215G

K216S

D229S

30

These single mutations can be engineered into a G $\alpha$ S backbone singly and in pairwise combinations. In addition, they can be introduced in the context of both the full length G $\alpha$ S and the GPA<sub>41</sub>-G $\alpha$ S hybrid described previously. All are predicted to improve the coupling of G $\alpha$  subunits to yeast G $\beta\gamma$  subunits by virtue of improved electrostatic and hydrophobic contacts between this region and the regions of G $\beta$  defined by Whiteway and co-workers (Whiteway et al (*supra*) that define site(s) that interact with GPA1).

35

In summary, the identification of hybrid G $\alpha$  subunits that couple to the yeast pheromone pathway has led to the following general observations. First, GPABAMH1

5 hybrids associate with yeast  $G\beta\gamma$ , therefore at a minimum these hybrids contain the determinants in GPA1 necessary for coupling to the pheromone response pathway. Second, the amino terminal 41 residues of GPA1 contain sufficient determinants to facilitate coupling of  $G\alpha$  hybrids to yeast  $G\beta\gamma$  in some, but not all, instances, and that  
10 some  $G\alpha$  subunits contain regions outside of the first 41 residues that are sufficiently similar to those in GPA1 to facilitate interaction with GPA1 even in the absence of the amino terminal 41 residues of GPA1. Third, there are other determinants in the first 310 residues of GPA1 that are involved in coupling  $G\alpha$  subunits to yeast  $G\beta\gamma$  subunits.

The various classes of hybrids noted above are not mutually exclusive. For example, a GPA1 containing GPA1-41 could also feature the L203K mutation.

15 While, for the sake of simplicity, hybrids of yeast GPA1 and a mammalian  $G\alpha_s$  have been described here, it will be appreciated that hybrids may be made of other yeast  $G\alpha$  subunits and/or other mammalian  $G\alpha$  subunits, notably mammalian  $G\alpha_i$  subunits. Moreover, while the described hybrids are constructed from two parental proteins, hybrids of three or more parental proteins are also possible.

20 As shown in the Examples, chimeric  $G\alpha$  subunits have been especially useful in coupling receptors to  $G\alpha_i$  species.

#### *iv. Expression of $G\alpha$*

Kang et al. *supra* reported that several classes of native mammalian  $G\alpha$  subunits  
25 were able to interact functionally with yeast  $\alpha$  subunits when expression of  $G\alpha$  was driven from a constitutively active, strong promoter (PGK) or from a strong inducible promoter (CUP). These authors reported that rat  $G\alpha_s$ ,  $G\alpha_i2$  or  $G\alpha_o$  expressed at high level coupled to yeast  $\beta\gamma$ . High level expression of mammalian  $G\alpha$  (i.e. non-stoichiometric with respect to yeast  $\beta\gamma$ ) is not preferred for uses like those described in  
30 this application. Reconstruction of G protein-coupled receptor signal transduction in yeast requires the signaling component of the heterotrimeric complex ( $G\beta\gamma$ ) to be present stoichiometrically with  $G\alpha$  subunits. An excess of  $G\alpha$  subunits (as was required for coupling of mammalian  $G\alpha_i2$  and  $G\alpha_o$  to yeast  $G\beta\gamma$  in Kang et al.) would dampen the signal in systems where  $G\beta\gamma$  subunits transduce the signal. An excess of  $G\alpha$   
35 subunits raises the background level of signaling in the system. Preferably, levels of  $G\alpha$  and  $G\beta\gamma$  subunits are balanced. For example, heterologous  $G\alpha$  subunits may be expressed from a low copy (CEN ARS) vector containing the endogenous yeast GPA1 promoter and the GPA1 3' untranslated region. The minimum criterion, applied to a heterologous  $G\alpha$  subunit with respect to its ability to couple functionally to the yeast  
40 pheromone pathway, is that it complement a *gpa1* genotype when expressed from the

5 GPAI promoter on low copy plasmids or from an integrated, single copy gene. In the work described in this application, heterologous  $G\alpha$  subunits have been assayed in two biological systems. In the first assay heterologous  $G\alpha$  subunits are tested for an ability to functionally complement the growth arrest phenotype of *gpaI* strains. In the second assay the transcription of a *fusI-HIS3* reporter gene is used to measure the extent to which the pheromone response pathway is activated, and hence the extent to which the heterologous  $G\alpha$  subunit sequesters the endogenous yeast  $G\beta\gamma$  complex. Mammalian  $G\alpha_s$ ,  $G\alpha_i2$ ,  $G\alpha_i3$ ,  $G\alpha_q$ ,  $G\alpha_{11}$ ,  $G\alpha_{16}$ ,  $G\alpha_{o_a}$ ,  $G\alpha_{o_b}$ , and  $G\alpha_z$  from rat, murine or human origins were expressed from a low copy, CEN ARS vector containing the GPAI promoter. Functional complementation of *gpaI* strains was not observed in either assay system with any of these full-length  $G\alpha$  constructs with the exception of rat and human  $G\alpha_S$ .

#### D. Chimeric Yeast $\beta\gamma$ subunits

In addition to or in place of modifying G protein  $G\alpha$  subunits, yeast or heterologous  $G\beta$  or  $G\gamma$  subunits can be modified. The methods described above with regard to  $G\alpha$  modification can be used to alter either or both of these subunits as well. For example, alignments of the yeast sequence and heterologous sequences can be made and combined with information regarding important functional domains. Such information can then be used to provide guidance in making mutations in yeast or heterologous sequences. Likewise, chimeric  $G\beta$  or  $G\gamma$  molecules can be constructed to enhance the coupling of heterologous GPCRs to a yeast pheromone signaling pathway.

The yeast STE4 and STE18 are related to the metazoan G protein  $\beta$  and  $\gamma$  subunits, respectively (Whiteway et al. 1989. Cell. 56:467). The  $\beta$  and  $\gamma$  subunits must be capable of interaction with one another as well as with the  $\alpha$  subunit and with the effector. Previous work has suggested that mammalian  $\beta$  or  $\gamma$  subunits are divergent enough from their yeast homologues that they cannot functionally replace STE4 or STE 18. (Coria et al. 1996. Yeast. 12:41). Thus, in preferred embodiments, modifications are made to heterologous  $G\beta$  or  $G\gamma$  subunits expressed in yeast and/or chimeric subunits are made to enhance heterologous receptor coupling.

The primary structure of G-protein  $\beta$  subunits is highly conserved from yeast to humans; Ste4 shares approximately 40% identity with human  $G\beta$  isoforms (Leberer et al. 1992 EMBO Journal 11:4085). STE 4 and the  $G\beta$ s are 420, and 340 or 341 amino acids long, respectively, and belong to the family of proteins with WD-40 motifs (van der Voorn and Ploegh. 1992. FEBS Lett. 307:131). These motifs can be used to divide  $G\beta$  and STE4 into eight blocks (Coria et al. Yeast 1996. 12:41). Among the mammalian  $G\beta$ s, some have been found to exhibit  $G\gamma$  subunit selectivity (Pronin and

5 Gautham. 1992. *Proc. Natl. Acad. Sci. USA* 89:6220; Schmidt et al. 1992. *J. Biol. Chem.* 267:13807; Kleuss et al. 1992. *Nature.* 358:424). An alignment of the metazoan and yeast G protein  $\beta$  subunits is provided by Corai et al. (1996. *Yeast.* 12:41). Such an alignment can be used to provide guidance for making mutations to G protein  $\beta$  subunits as described for  $G\alpha$  above. In addition, certain regions of STE4 have  
10 been found to be important and thus, may be less amenable to manipulation than other portions of the polypeptide. For example, the c-terminus of the STE4 product is essential for downstream signaling (Coria et al. 1995. *FEBS Letters* 367:122). Mutations to two small regions in the amino terminal half of Ste4 have also been shown to inhibit signaling (Leberer et al. *supra*). Mutations which influence the interaction of  
15 STE4 and GPA1 have also been identified; mutations to the second copy of the WD40 repeat can be modified to reduce the interaction between STE4 and GPA1, without influencing other aspects of STE4 function (Whiteway et al. 1994. *Mol. Cell. Biol.* 14:3223)

The Gys, including STE18, diverge more strongly from each other than do the  
20  $G\beta$ s. Even among the mammalian G protein  $\gamma$  subunits, there is a fair amount of divergence. The  $\gamma$  subunit may determine the functional specificity of the  $\beta\gamma$  subunit complex. Complete cDNAs for the  $\gamma 1$  subunit from bovine retina (Hurley et al. *Proc. Nat'l Acad. Sci. USA.* 1984. 81:6948) the  $\gamma 1$ ,  $\gamma 3$ , and  $\gamma 7$  subunits from bovine brain (Robishaw et al. *J. Biol. Chem.* 1989. 264:15758; Gautam et al. *Science.* 1989.  
25 244:971; Gautam et al. *Proc. Nat'l Acad. Sci. USA.* 1990 87:7973; Cali et al. *J. Biol. Chem.* 1992. 267:24023), and the  $\gamma 5$  subunit from bovine and rat liver (Gisher et al., 1992. 12:1585) have been reported.

The STE18 gene of yeast terminates with a CAAX box (where A is an aliphatic amino acid, and X is any uncharged amino acid). This sequence is involved in  
30 prenylation of  $G\gamma$  and is likely important in the localization of  $G\gamma$  to the membrane and may, thus, be less amenable to manipulation than other portions of the sequence. (Kurjan. 1992. *Ann. Rev. Biochem.* 61:1097). Saturation mutagenesis has also provided insight into regions of STE18 that are important in STE18 function. Mutations in STE18 which compensate for mutations in STE4 were identified at serine 65,  
35 threonine 71, and valine 80. Dominant negative alleles of the STE18 gene were also identified (Whiteway et al. 1992. *Biochem. Cell. Biol.* 70:1230). These truncated proteins were found to lack the carboxyl terminus of STE 18, including the CAAX box (Whiteway et al. *supra*).

An alignment of yeast  $G\gamma$ , STE18, and mammalian Gys can be made as indicated  
40 for the other G protein subunits. Such an alignment can be used in constructing mutant

- 5 Gy subunits or chimeric Gy subunits. In preferred embodiments, mammalian Gy2 is used in making G protein  $\gamma$  subunit chimeras.

### VIII. Leader Sequences

It has been demonstrated that most of the mammalian extracellular, secreted proteins are poorly secreted when expressed in yeast. However, in many cases their secretion levels are markedly increased when their native signal sequences are replaced by the signal sequences of yeast proteins that interact more efficiently with the ER translocation complex. Specifically, the signal sequences of yeast invertase and acid phosphatase have been widely used in biotechnology to direct the secretory expression of the heterologous proteins. However, it is well established that even though many foreign proteins are targeted to the ER by the yeast signal sequences, not all of them advance further in the secretory pathway. The major problem appears to be in the misfolding and/or improper glycosylation of the heterologous proteins that results in their retention in the ER by the quality control apparatus of the yeast cell.

In many cases, the leader sequence of a precursor of yeast mating pheromone,  $\alpha$ -factor, has been used successfully to overcome this problem (Brake, A.J. (1989) in *Yeast Genetic Engineering* (Barr, P.J., Brake, A.J., and Valenzuela, P., eds) pp. 269-280, Butterworths, London; Brake, A.J. (1990) *Meth. Enzymol.* 185, 408-441., and references cited therein). This sequence, in addition to the N-terminal signal peptide of 17 residues, includes a hydrophilic pro-region which contains 72 residues and bears three sites of N-linked glycosylation. The pro-region is extensively glycosylated in the ER and Golgi and is cleaved by Kex2 endopeptidase in the late Golgi compartment. The presence of the pro-region at the N-terminus has been demonstrated to promote transport of heterologous proteins from the ER to the periplasm. It is likely that the pro-region can somehow facilitate correct protein folding. Alternatively, it may be recognized by the quality control apparatus as a properly folded structural unit thus allowing an entire fusion protein to leave the ER.

The invertase leader can also be used. This leader sequence has been demonstrated to be cleaved from nascent invertase peptide, or nascent heterologous peptide, in the course of translocation into the endoplasmic reticulum.

10

### *A. Peptide Expression*

In certain embodiments, such a leader sequence can be used to express a peptide library of the present invention. Yeast cells are bounded by a lipid bilayer called the plasma membrane. Between this plasma membrane and the cell wall is the periplasmic

5 space. Peptides secreted by yeast cells cross the plasma membrane through a variety of mechanisms and thereby enter the periplasmic space. The secreted peptides are then free to interact with other molecules that are present in the periplasm or displayed on the outer surface of the plasma membrane. The peptides may either undergo re-uptake into the cell, transit through the cell wall into the medium, or become degraded within the  
10 periplasmic space.

The test polypeptide library may be secreted into the periplasm by any of a number of exemplary mechanisms, depending on the nature of the expression system to which they are linked. In one embodiment, the peptide may be structurally linked to a yeast signal sequence, such as that present in the  $\alpha$ -factor precursor, which directs  
15 secretion through the endoplasmic reticulum and Golgi apparatus. Since this is the same route that the receptor protein follows in its journey to the plasma membrane, opportunity exists in cells expressing both the receptor and the peptide library for a specific peptide to interact with the receptor during transit through the secretory pathway. This has been postulated to occur in mammalian cells exhibiting autocrine  
20 activation. Such interaction could yield activation of the response pathway during transit, which would still allow identification of those cells expressing a peptide agonist. For situations in which peptide antagonists to externally applied receptor agonist are sought, this system would still be effective, since both the peptide antagonist and receptor would be delivered to the outside of the cell in concert. Thus, those cells  
25 producing an antagonist would be selectable, since the peptide antagonist would be properly and timely situated to prevent the receptor from being stimulated by the externally applied agonist.

An alternative mechanism for delivering peptides to the periplasmic space is to use the ATP-dependent transporters of the STE6/MDR1 class. This transport pathway  
30 and the signals that direct a protein or peptide to this pathway are not as well characterized as is the endoplasmic reticulum-based secretory pathway. Nonetheless, these transporters apparently can efficiently export certain peptides directly across the plasma membrane, without the peptides having to transit the ER/Golgi pathway. It is anticipated that at least a subset of peptides can be secreted through this pathway by  
35 expressing the library in context of the  $\alpha$ -factor prosequence and terminal tetrapeptide. The possible advantage of this system is that the receptor and peptide do not come into contact until both are delivered to the external surface of the cell. Thus, this system strictly mimics the situation of an agonist or antagonist that is normally delivered from outside the cell. Use of either of the described pathways is within the scope of the  
40 invention.

5           The present invention does not require periplasmic secretion of peptides, or, if such secretion is provided, any particular secretion signal or transport pathway. In certain embodiments, peptides expressed with a signal sequence may bind to and activate receptors prior to their transport to the cell surface.

10           *B. Receptor Expression*

          In other embodiments, a leader sequence of a yeast secreted protein can be used to direct transport of heterologous receptors to the plasma membrane. Previous work has demonstrated the expression of foreign, secreted proteins in yeast cells using the  $\alpha$ -factor leader. However, when a heterologous membrane bound receptor, the rat M5  
15   receptor, was expressed using such a system, it was found that the heterologous GPCR did not functionally integrate into the yeast cell signaling pathway (Huang et al. *Biochem. and Biophys. Res. Comm.* 1992. 182:1180). The transport of both secreted and transmembrane proteins into the endoplasmic reticulum in yeast is promoted by the same protein translocation complex, including the Sec61, Sec62 and Sec63 proteins. All  
20   the secreted proteins possess a signal sequence at their N-termini which is recognized by the translocation complex and serves as an ER targeting signal. A typical signal sequence is comprised of several positively charged residues at the N-terminus followed by a hydrophobic core and a C-terminal site of processing by signal peptidase. Some transmembrane proteins, for example, metabotropic glutamate receptors and vasoactive  
25   intestinal polypeptide receptors, also possess the N-terminal signal sequences, whereas some do not. In the latter case, a first transmembrane domain is believed to interact with the ER translocation machinery. The use of the  $\alpha$ -factor leader sequence may, therefore, be particularly desirable for functional expression of certain receptors.

30           In certain embodiments, it will be desirable to further modify the yeast cells of the present invention. For example, in one embodiment it will be desirable to disrupt the yeast calnexin-like gene, *CNE1*, to improve receptor transport from the endoplasmic reticulum to the Golgi. In yet other embodiments, it will be desirable to overexpress the gene encoding Ast1, to increase transport from the Golgi to the plasma membrane. In  
35   yet other embodiments, it will be desirable to disrupt *END3* and/or *END4*, to inhibit or prevent receptor internalization. Additionally or alternatively, the *CHC1* gene (clathrin-encoding) can be disrupted to inhibit or prevent receptor internalization. In another embodiment, it may be desirable to disrupt the *MVP-1* gene to inhibit or prevent transport from the Golgi to the prevacuolar compartment.

5     IX. Test Compounds

*Exogenously added compounds*

          A recent trend in medicinal chemistry includes the production of mixtures of compounds, referred to as libraries. While the use of libraries of peptides is well established in the art, new techniques have been developed which have allowed the  
10     production of mixtures of other compounds, such as benzodiazepines (Bunin et al. 1992. J. Am. Chem. Soc. 114:10987; DeWitt et al. 1993. Proc. Natl. Acad. Sci. USA 90:6909) peptoids (Zuckermann. 1994. J. Med. Chem. 37:2678) oligocarbamates (Cho et al. 1993. Science. 261:1303), and hydantoins (DeWitt et al. supra). Rebek et al. have described an approach for the synthesis of molecular libraries of small organic  
15     molecules with a diversity of 10<sup>4</sup>-10<sup>5</sup> (Carell et al. 1994. Angew. Chem. Int. Ed. Engl. 33:2059; Carell et al. Angew. Chem. Int. Ed. Engl. 1994. 33:2061).

          The compounds of the present invention can be obtained using any of the numerous approaches in combinatorial library methods known in the art, including: biological libraries; spatially addressable parallel solid phase or solution phase libraries,  
20     synthetic library methods requiring deconvolution, the 'one-bead one-compound' library method, and synthetic library methods using affinity chromatography selection. The biological library approach is limited to peptide libraries, while the other four approaches are applicable to peptide, non-peptide oligomer or small molecule libraries of compounds (Lam, K.S. Anticancer Drug Des. 1997. 12:145).

25     In one embodiment, the test compound is a peptide or peptidomimetic. In another, preferred embodiment, the compounds are small, organic non-peptidic compounds.

          Other exemplary methods for the synthesis of molecular libraries can be found in the art, for example in: Erb et al. 1994. Proc. Natl. Acad. Sci. USA 91:11422; Horwell  
30     et al. 1996 Immunopharmacology 33:68; and in Gallop et al. 1994. J. Med. Chem. 37:1233. In addition, libraries such as those described in the commonly owned applications U.S.S.N. 08/864,241, U.S.S.N. 08/864,240 and U.S.S.N. 08/835,623 can be used to provide compounds for testing in the present invention. The contents of each of these applications is expressly incorporated herein by this reference.

35     Libraries of compounds may be presented in solution (e.g., Houghten (1992) *Biotechniques* 13:412-421), or on beads (Lam (1991) *Nature* 354:82-84), chips (Fodor (1993) *Nature* 364:555-556), bacteria (Ladner USP 5,223,409), spores (Ladner USP '409), plasmids (Cull et al. (1992) *Proc Natl Acad Sci USA* 89:1865-1869) or on phage (Scott and Smith (1990) *Science* 249:386-390); (Devlin (1990) *Science* 249:404-406);



- 5 (Cwirla et al. (1990) *Proc. Natl. Acad. Sci.* 87:6378-6382); (Felici (1991) *J. Mol. Biol.* 222:301-310); (Ladner *supra.*).

In certain embodiments, the test compounds are exogenously added to the host cells expressing a recombinant receptor and compounds that modulate signal transduction via the receptor are selected. In other embodiments, the host cells express  
10 the compounds to be tested. For example, a culture of the subject host cells can be further modified to collectively express a peptide library as described in more detail in PCT Publication WO 94/23025 the contents of which is expressly incorporated herein by this reference.

Other types of peptide libraries may also be expressed, see, for example, U.S.  
15 Patents 5,270,181 and 5,292,646; and PCT publication WO94/ 02502). In still another embodiment, the combinatorial polypeptides are produced from a cDNA library.

Exemplary compounds which can be screened for activity include, but are not limited to, peptides, nucleic acids, carbohydrates, small organic molecules, and natural product extract libraries. In such embodiments, both compounds which agonize or  
20 antagonize the receptor- or channel-mediated signaling function can be selected and identified.

#### *Autocrine host Cells*

In certain embodiments, host cells, e.g., can be engineered to produce the  
25 compounds to be tested. This assay system has the advantage of increasing the effective concentration of the compound to be tested. In one embodiment, a method such as that described in WO 94/23025 can be utilized.

Other methods can also be used. For example, peptide libraries are systems which simultaneously display, in a form which permits interaction with a target, a highly  
30 diverse and numerous collection of peptides. These peptides may be presented in solution (Houghten (1992) *Biotechniques* 13:412-421), or on beads (Lam (1991) *Nature* 354:82-84), chips (Fodor (1993) *Nature* 364:555-556), bacteria (Ladner USP 5,223,409), spores (Ladner USP '409), plasmids (Cull et al. (1992) *Proc Natl Acad Sci USA* 89:1865-1869) or on phage (Scott and Smith (1990) *Science* 249:386-390); (Devlin  
35 (1990) *Science* 249:404-406); (Cwirla et al. (1990) *Proc. Natl. Acad. Sci.* 87:6378-6382); (Felici (1991) *J. Mol. Biol.* 222:301-310); (Ladner *supra.*). Many of these systems are limited in terms of the maximum length of the peptide or the composition of the peptide (e.g., Cys excluded). Steric factors, such as the proximity of a support, may interfere with binding. Usually, the screening is for binding in vitro to an artificially  
40 presented target, not for activation or inhibition of a cellular signal transduction pathway

5 in a living cell. While a cell surface receptor may be used as a target, the screening will not reveal whether the binding of the peptide caused an allosteric change in the conformation of the receptor.

The Ladner et al. patent, USSN 5,096,815, describes a method of identifying novel proteins or polypeptides with a desired DNA binding activity. Semi-random  
10 ("variegated") DNA encoding a large number of different potential binding proteins is introduced, in expressible form, into suitable yeast cells. The target DNA sequence is incorporated into a genetically engineered operon such that the binding of the protein or polypeptide will prevent expression of a gene product that is deleterious to the gene under selective conditions. Cells which survive the selective conditions are thus cells  
15 which express a protein which binds the target DNA. While it is taught that yeast cells may be used for testing, bacterial cells are preferred. The interactions between the protein and the target DNA occur only in the cell (and then only in the nucleus), not in the periplasm or cytoplasm, and the target is a nucleic acid, and not a receptor protein. Substitution of random peptide sequences for functional domains in cellular proteins  
20 permits some determination of the specific sequence requirements for the accomplishment of function. Though the details of the recognition phenomena which operate in the localization of proteins within cells remain largely unknown, the constraints on sequence variation of mitochondrial targeting sequences and protein secretion signal sequences have been elucidated using random peptides (Lemire et al., *J. Biol. Chem.*(1989) 264, 20206 and Kaiser et al. (1987) *Science* 235:312, respectively).  
25

In certain embodiments of the instant invention, the compounds tested are in the form of peptides from a peptide library. The peptide library of the present invention takes the form of a cell culture, in which essentially each cell expresses one, and usually only one, peptide of the library. While the diversity of the library is maximized if each  
30 cell produces a peptide of a different sequence, it is usually prudent to construct the library so there is some redundancy. Depending on size, the combinatorial peptides of the library can be expressed as is, or can be incorporated into larger fusion proteins. The fusion protein can provide, for example, stability against degradation or denaturation, as well as a secretion signal if secreted. In an exemplary embodiment of a library for  
35 intracellular expression, e.g., for use in conjunction with intracellular target receptors, the polypeptide library is expressed as thioredoxin fusion proteins (see, for example, U.S. Patents 5,270,181 and 5,292,646; and PCT publication WO94/ 02502). The combinatorial peptide can be attached on the terminus of the thioredoxin protein, or, for short peptide libraries, inserted into the so-called active loop.

40 In one embodiment, the peptide library is derived to express a combinatorial library of polypeptides which are not based on any known sequence, nor derived from

5 cDNA. That is, the sequences of the library are largely random. In preferred  
embodiments, the combinatorial polypeptides are in the range of 3-100 amino acids in  
length, more preferably at least 5-50, and even more preferably at least 10, 13, 15, 20 or  
25 amino acid residues in length. Preferably, the polypeptides of the library are of  
uniform length. It will be understood that the length of the combinatorial peptide does  
10 not reflect any extraneous sequences which may be present in order to facilitate  
expression, e.g., such as signal sequences or invariant portions of a fusion protein.

In another embodiment, the peptide library is a combinatorial library of  
polypeptides which are based at least in part on a known polypeptide sequence or a  
portion thereof (not a cDNA library). That is, the sequences of the library is semi-  
15 random, being derived by combinatorial mutagenesis of a known sequence. See, for  
example, Ladner et al. PCT publication WO 90/02909; Garrard et al., PCT publication  
WO 92/09690; Marks et al. (1992) *J. Biol. Chem.* 267:16007-16010; Griffiths et al.  
(1993) *EMBO J* 12:725-734; Clackson et al. (1991) *Nature* 352:624-628; and Barbas et  
al. (1992) *PNAS* 89:4457-4461. Accordingly, polypeptide(s) which are known ligands  
20 for a target receptor can be mutagenized by standard techniques to derive a variegated  
library of polypeptide sequences which can further be screened for agonists and/or  
antagonists. For example, the DNA encoding a surrogate ligand can be mutagenized to  
generate a library encoding peptides with some relationship to the original peptide. This  
library can be expressed in a reagent cell of the present invention, and other receptor  
25 activators can be isolated from the library. This may permit the identification of even  
more potent surrogate ligands.

In still another embodiment, the combinatorial polypeptides are produced from a  
cDNA library.

In a preferred embodiment of the present invention, the host cells collectively  
30 produce a "peptide library", preferably including at least  $10^3$  to  $10^7$  different peptides, so  
that diverse peptides may be simultaneously assayed for the ability to interact with the  
exogenous receptor. In an especially preferred embodiment, at least some peptides of  
the peptide library are secreted into the periplasm, where they may interact with the  
"extracellular" binding site(s) of an exogenous receptor. They thus mimic more closely  
35 the clinical interaction of drugs with cellular receptors. This embodiment optionally  
may be further improved (in assays not requiring pheromone secretion) by preventing  
pheromone secretion, and thereby avoiding competition between the peptide and the  
pheromone for signal peptidase and other components of the secretion system.

In certain embodiments of the present invention, the peptides of the library are  
40 encoded by a mixture of DNA molecules of different sequence. Each peptide-encoding  
DNA molecule is ligated with a vector DNA molecule and the resulting recombinant

5 DNA molecule is introduced into a host cell. Since it is a matter of chance which peptide-encoding DNA molecule is introduced into a particular cell, it is not predictable which peptide that cell will produce. However, based on a knowledge of the manner in which the mixture was prepared, one may make certain statistical predictions about the mixture of peptides in the peptide library.

10 The peptides of the library can be composed of constant and variable residues. If the *n*th residue is the same for all peptides of the library, it is said to be constant. If the *n*th residue varies, depending on the peptide in question, the residue is a variable one. The peptides of the library will have at least one, and usually more than one, variable residue. A variable residue may vary among any of two to all twenty of the genetically  
15 encoded amino acids; the variable residues of the peptide may vary in the same or different manner. Moreover, the frequency of occurrence of the allowed amino acids at a particular residue position may be the same or different. The peptide may also have one or more constant residues.

There are two principal ways in which to prepare the required DNA mixture. In  
20 one method, the DNAs are synthesized a base at a time. When variation is desired, at a base position dictated by the Genetic Code, a suitable mixture of nucleotides is reacted with the nascent DNA, rather than the pure nucleotide reagent of conventional polynucleotide synthesis.

The second method provides more exact control over the amino acid variation.  
25 First, trinucleotide reagents are prepared, each trinucleotide being a codon of one (and only one) of the amino acids to be featured in the peptide library. When a particular variable residue is to be synthesized, a mixture is made of the appropriate trinucleotides and reacted with the nascent DNA. Once the necessary "degenerate" DNA is complete, it must be joined with the DNA sequences necessary to assure the expression of the  
30 peptide, as discussed in more detail below, and the complete DNA construct must be introduced into the yeast cell.

#### X. Screening and Selection: Assays of Second Messenger Generation

When screening for bioactivity of compounds, intracellular second messenger  
35 generation can be measured directly. A variety of intracellular effectors have been identified as being G-protein-regulated, including adenylyl cyclase, cyclic GMP, phosphodiesterases, phosphoinositidase C, and phospholipase A<sub>2</sub>. In addition, G proteins interact with a range of ion channels and are able to inhibit certain voltage-sensitive Ca<sup>++</sup> transients, as well as stimulating cardiac K<sup>+</sup> channels.

5           In one embodiment, the GTPase enzymatic activity by G proteins can be measured in plasma membrane preparations by determining the breakdown of  $\gamma^{32}\text{P}$  GTP using techniques that are known in the art (For example, see *Signal Transduction: A Practical Approach*. G. Milligan, Ed. Oxford University Press, Oxford England). When receptors that modulate cAMP are tested, it will be possible to use standard  
10 techniques for cAMP detection, such as competitive assays which quantitate [ $^3\text{H}$ ]cAMP in the presence of unlabelled cAMP.

Certain receptors stimulate the activity of phospholipase C which stimulates the breakdown of phosphatidylinositol 4,5, biphosphate to 1,4,5-IP<sub>3</sub> (which mobilizes intracellular  $\text{Ca}^{++}$ ) and diacylglycerol (DAG) (which activates protein kinase C).  
15 Inositol lipids can be extracted and analyzed using standard lipid extraction techniques. DAG can also be measured using thin-layer chromatography. Water soluble derivatives of all three inositol lipids (IP<sub>1</sub>, IP<sub>2</sub>, IP<sub>3</sub>) can also be quantitated using radiolabelling techniques or HPLC.

The mobilization of intracellular calcium or the influx of calcium from outside  
20 the cell can be measured using standard techniques. The choice of the appropriate calcium indicator, fluorescent, bioluminescent, metallochromic, or  $\text{Ca}^{++}$ -sensitive microelectrodes depends on the cell type and the magnitude and time constant of the event under study (Borle (1990) *Environ Health Perspect* 84:45-56). As an exemplary method of  $\text{Ca}^{++}$  detection, cells could be loaded with the  $\text{Ca}^{++}$ -sensitive fluorescent dye  
25 fura-2 or indo-1, using standard methods, and any change in  $\text{Ca}^{++}$  measured using a fluorometer.

The other product of PIP<sub>2</sub> breakdown, DAG can also be produced from phosphatidyl choline. The breakdown of this phospholipid in response to receptor-mediated signaling can also be measured using a variety of radiolabelling techniques.

30           The activation of phospholipase A<sub>2</sub> can easily be quantitated using known techniques, including, for example, the generation of arachadonate in the cell.

In the case of certain receptors, it may be desirable to screen for changes in cellular phosphorylation. Such assay formats may be useful when the receptor of interest is a receptor tyrosine kinase. For example, yeast transformed with the FGF  
35 receptor and a ligand which binds the FGF receptor could be screened using colony immunoblotting (Lyons and Nelson (1984) *Proc. Natl. Acad. Sci. USA* 81:7426-7430) using anti-phosphotyrosine. In addition, tests for phosphorylation could be useful when a receptor which may not itself be a tyrosine kinase, activates protein kinases that function downstream in the signal transduction pathway. Likewise, it is noted that  
40 protein phosphorylation also plays a critical role in cascades that serve to amplify signals generated at the receptor. Multi-kinase cascades allow not only signal

5 amplification but also signal divergence to multiple effectors that are often cell-type specific, allowing a growth factor to stimulate mitosis of one cell and differentiation of another.

One such cascade is the MAP kinase pathway that appears to mediate both mitogenic, differentiation and stress responses in different cell types. Stimulation of growth factor receptors results in Ras activation followed by the sequential activation of c-Raf, MEK, and p44 and p42 MAP kinases (ERK1 and ERK2). Activated MAP kinase then phosphorylates many key regulatory proteins, including p90RSK and Elk-1 that are phosphorylated when MAP kinase translocates to the nucleus. Homologous pathways exist in mammalian and yeast cells. For instance, an essential part of the *S. cerevisiae* pheromone signaling pathway is comprised of a protein kinase cascade composed of the products of the STE11, STE7, and FUS3/KSS1 genes (the latter pair are distinct and functionally redundant). Accordingly, phosphorylation and/or activation of members of this kinase cascade can be detected and used to quantitate receptor engagement. Phosphotyrosine specific antibodies are available to measure increases in tyrosine phosphorylation and phospho-specific antibodies are commercially available (New England Biolabs, Beverly, MA).

Modified methods for detecting receptor-mediated signal transduction exist and one of skill in the art will recognize suitable methods that may be used to substitute for the example methods listed.

25 In one embodiment, the indicator gene can be used for detection. In one embodiment an indicator gene is an unmodified endogenous gene. For example, the instant method can rely on detecting the transcriptional level of such pheromone response pathway responsive endogenous genes as the *Bar1* or *Fus1*, *Fus 2*, mating factor, *Ste3* *Ste13*, *Kex1*, *Ste2*, *Ste6*, *Ste7*, *Sst2*, or *Chs1*. (Appletauer and Zchstetter. 1989. Eur. J. Biochem. 181:243)

In other embodiments, the sensitivity of an endogenous indicator gene can be enhanced by manipulating the promoter sequence at the natural locus for the indicator gene. Such manipulation may range from point mutations to the endogenous regulatory elements to gross replacement of all or substantial portions of the regulatory elements. The previous discussion of mutations with regard to G proteins and G protein-coupled receptors is reiterated here.

For example, in the case of the *Bar1* gene, the promoter of the gene can be modified to enhance the transcription of *Bar1* upon activation of the yeast pheromone response pathway. *Bar1* gene transcription is inactivated upon exposure of yeast cells to mating factor. The sequence of the *Bar1* gene is known in the art (see e.g., U.S. patent 4,613,572). Moreover, the sequences required for  $\alpha$ -factor-enhanced expression of the

5 Bar1, and other pheromone responsive genes have been identified. (Appeltauer and Achstetter 1989. Eur. J. Biochem. 181:243; Hagen et al. 1991. Mol. Cell. Biol. 11:2952). In an exemplary embodiment, the yeast *Bar1* promoter can be engineered by mutagenesis to be more responsive, e.g., to more strongly promoter gene transcription, upon stimulation of the yeast pheromone pathway. Standard techniques for  
10 mutagenizing the promoter can be used. In such embodiments, it is desirable that the conserved oligonucleotide motif described by Appeltaure et al. be conserved.

In yet other embodiments, rather than measuring second messenger production or alterations in transcription, the activity of endogenous yeast proteins can be assayed. For example, in one embodiment, the signal transduction pathway of the receptor  
15 upregulates expression or otherwise activates an enzyme which is capable of modifying a substrate which can be added to the cell. The signal can be detected by using a detectable substrate, in which case loss of the substrate signal is monitored, or alternatively, by using a substrate which produces a detectable product. In certain embodiments, the substrate is naturally occurring. Alternatively, the substrate can be  
20 non-naturally occurring. In preferred embodiments, BAR1 activity can be measured.

In other embodiments, the modulation of a receptor by a test compound can result in a change in the transcription of a gene, which is not normally pheromone responsive. In preferred embodiments, the gene is easily detectable. For example, in a preferred embodiment, the subject assay can be used to measure Pho5, a secreted acid  
25 phosphatase. Acid phosphatase activity can be measured using standard techniques.

In other embodiments, reporter gene constructs can be used. Reporter gene constructs are prepared by operatively linking a reporter gene with at least one transcriptional regulatory element. If only one transcriptional regulatory element is included it must be a regulatable promoter. At least one of the selected transcriptional  
30 regulatory elements must be indirectly or directly regulated by the activity of the selected cell-surface receptor whereby activity of the receptor can be monitored via transcription of the reporter genes.

Many reporter genes and transcriptional regulatory elements are known to those of skill in the art and others may be identified or synthesized by methods known to those  
35 of skill in the art. Reporter genes include any gene that expresses a detectable gene product, which may be RNA or protein. Preferred reporter genes are those that are readily detectable. The reporter gene may also be included in the construct in the form of a fusion gene with a gene that includes desired transcriptional regulatory sequences or exhibits other desirable properties.

40 Examples of reporter genes include, but are not limited to CAT (chloramphenicol acetyl transferase) (Alton and Vapnek (1979), Nature 282: 864-869) luciferase, and other

5 enzyme detection systems, such as beta-galactosidase; firefly luciferase (deWet et al. (1987), *Mol. Cell. Biol.* 7:725-737); bacterial luciferase (Engebrecht and Silverman (1984), *PNAS* 1: 4154-4158; Baldwin et al. (1984), *Biochemistry* 23: 3663-3667); alkaline phosphatase (Toh et al. (1989) *Eur. J. Biochem.* 182: 231-238, Hall et al. (1983) *J. Mol. Appl. Gen.* 2: 101), human placental secreted alkaline phosphatase (Cullen and  
10 Malim (1992) *Methods in Enzymol.* 216:362-368) and green fluorescent protein (U.S. patent 5,491,084; WO96/23898).

Transcriptional control elements include, but are not limited to, promoters, enhancers, and repressor and activator binding sites. Suitable transcriptional regulatory elements may be derived from the transcriptional regulatory regions of genes whose  
15 expression is rapidly induced, generally within minutes, of contact between the cell surface protein and the effector protein that modulates the activity of the cell surface protein. Examples of such genes include, but are not limited to, the immediate early genes (see, Sheng et al. (1990) *Neuron* 4: 477-485), such as c-fos. Immediate early genes are genes that are rapidly induced upon binding of a ligand to a cell surface  
20 protein. The transcriptional control elements that are preferred for use in the gene constructs include transcriptional control elements from immediate early genes, elements derived from other genes that exhibit some or all of the characteristics of the immediate early genes, or synthetic elements that are constructed such that genes in operative linkage therewith exhibit such characteristics. The characteristics of preferred genes  
25 from which the transcriptional control elements are derived include, but are not limited to, low or undetectable expression in quiescent cells, rapid induction at the transcriptional level within minutes of extracellular stimulation, induction that is transient and independent of new protein synthesis, subsequent shut-off of transcription requires new protein synthesis, and mRNAs transcribed from these genes have a short half-life.  
30 It is not necessary for all of these properties to be present.

Other promoters and transcriptional control elements, in addition to those described above, include the vasoactive intestinal peptide (VIP) gene promoter (cAMP responsive; Fink et al. (1988), *Proc. Natl. Acad. Sci.* 85:6662-6666); the somatostatin gene promoter (cAMP responsive; Montminy et al. (1986), *Proc. Natl. Acad. Sci.*  
35 83:6682-6686); the proenkephalin promoter (responsive to cAMP, nicotinic agonists, and phorbol esters; Comb et al. (1986), *Nature* 323:353-356); the phosphoenolpyruvate carboxy-kinase gene promoter (cAMP responsive; Short et al. (1986), *J. Biol. Chem.* 261:9721-9726); the NGFI-A gene promoter (responsive to NGF, cAMP, and serum; Changelian et al. (1989). *Proc. Natl. Acad. Sci.* 86:377-381); and others that may be  
40 known to or prepared by those of skill in the art.



5           In certain assays it may be desirable to use changes in growth in the screening procedure. For example, one of the consequences of activation of the pheromone signal pathway in wild-type yeast is growth arrest. If one is testing for an antagonist of a G protein-coupled receptor, this normal response of growth arrest can be used to select cells in which the pheromone response pathway is inhibited. That is, cells exposed to  
10 both a known agonist and a peptide of unknown activity will be growth arrested if the peptide is neutral or an agonist, but will grow normally if the peptide is an antagonist. Thus, the growth arrest response can be used to advantage to discover peptides that function as antagonists.

          In certain embodiments, when searching for compounds which can function as  
15 agonists of G protein-coupled receptors, or other pheromone system proteins, the growth arrest consequent to activation of the pheromone response pathway is an undesirable effect since cells that bind agonists stop growing while surrounding cells that fail to bind agonists will continue to grow. The cells of interest, then, will be overgrown or their detection obscured by the background cells, confounding identification of the cells of  
20 interest. To overcome this problem the present invention teaches engineering the cell such that: 1) growth arrest does not occur as a result of exogenous signal pathway activation (e.g., by inactivating the FAR1 gene); and/or 2) a selective growth advantage is conferred by activating the pathway (e.g., by transforming an auxotrophic mutant with a HIS3 gene under the control of a pheromone-responsive promoter, and applying  
25 selective conditions).

          Alternatively, the promoter may be one which is repressed by the receptor pathway, thereby preventing expression of a product which is deleterious to the cell. With a receptor repressed promoter, one screens for agonists by linking the promoter to a deleterious gene, and for antagonists, by linking it to a beneficial gene. Repression  
30 may be achieved by operably linking a receptor- induced promoter to a gene encoding mRNA which is antisense to at least a portion of the mRNA encoded by the marker gene (whether in the coding or flanking regions), so as to inhibit translation of that mRNA. Repression may also be obtained by linking a receptor-induced promoter to a gene encoding a DNA-binding repressor protein, and incorporating a suitable operator site  
35 into the promoter or other suitable region of the marker gene.

          In the case of yeast, exemplary positively selectable (beneficial) genes include the following: *URA3*, *LYS2*, *HIS3*, *LEU2*, *TRP1*; *ADE1,2,3,4,5,7,8*; *ARG1*, 3, 4, 5, 6, 8; *HIS1*, 4, 5; *ILV1*, 2, 5; *THR1*, 4; *TRP2*, 3, 4, 5; *LEU1*, 4; *MET2,3,4,8,9,14,16,19*; *URA1,2,4,5,10*; *HOM3,6*; *ASP3*; *CHO1*; *ARO 2,7*; *CYS3*; *OLE1*; *INO1,2,4*; *PR01,3*  
40 Countless other genes are potential selective markers. The above are involved in well-characterized biosynthetic pathways. The imidazoleglycerol phosphate dehydratase

5 (IGP dehydratase) gene (HIS3) is preferred because it is both quite sensitive and can be selected over a broad range of expression levels. In the simplest case, the cell is auxotrophic for histidine (requires histidine for growth) in the absence of activation. Activation leads to synthesis of the enzyme and the cell becomes prototrophic for histidine (does not require histidine). Thus the selection is for growth in the absence of  
10 histidine. Since only a few molecules per cell of IGP dehydratase are required for histidine prototrophy, the assay is very sensitive.

In another version of the assay, cells can be selected for resistance to aminotriazole (AT), a drug that inhibits the activity of IGP dehydratase. Cells with low, fixed level of expression of HIS3 are sensitive to the drug, while cells with higher levels  
15 are resistant. The amount of AT can be selected to inhibit cells with a basal level of HIS3 expression (whatever that level is) but allow growth of cells with an induced level of expression. In this case selection is for growth in the absence of histidine and in the presence of a suitable level of AT.

In appropriate assays, so-called counterselectable or negatively selectable genes  
20 may be used. Suitable genes include: URA3 (orotidine-5'-phosphate decarboxylase; inhibits growth on 5-fluoroorotic acid), LYS2 (2-aminoadipate reductase; inhibits growth on  $\alpha$ -aminoadipate as sole nitrogen source), CYH2 (encodes ribosomal protein L29; cycloheximide-sensitive allele is dominant to resistant allele), CAN1 (encodes arginine permease; null allele confers resistance to the arginine analog canavanine), and  
25 other recessive drug-resistant markers.

In one example, the reporter gene effects yeast cell growth. The natural response to signal transduction via the yeast pheromone system response pathway is for cells to undergo growth arrest. This is a preferred way to select for antagonists of a ligand/receptor pair that stimulates a the pathway. An antagonist would inhibit the  
30 activation of the pathway; hence, the cell would be able to grow. Thus, the FAR1 gene may be considered an endogenous counterselectable marker. The FAR1 gene is preferably inactivated when screening for agonist activity.

The reporter gene may also be a screenable gene. The screened characteristic may be a change in cell morphology, metabolism or other screenable features. Suitable  
35 markers include beta-galactosidase (Xgal, C<sub>12</sub>FDG, Salmon-gal, Magenta-Gal (latter two from Biosynth Ag)), alkaline phosphatase, horseradish peroxidase, exo-glucanase (product of yeast exb1 gene; nonessential, secreted); luciferase; bacterial green fluorescent protein; (human placental) secreted alkaline phosphatase (SEAP); and chloramphenicol transferase (CAT). Some of the above can be engineered so that they  
40 are secreted (although not  $\beta$ -galactosidase). A preferred screenable marker gene is beta-

- 5 galactosidase; yeast cells expressing the enzyme convert the colorless substrate Xgal into a blue pigment. Again, the promoter may be receptor-induced or receptor-inhibited.

### XI. Other Optional Alterations to Yeast Cells

- 10 The choice of appropriate host cell will also be influenced by the choice of detection signal. For instance, reporter constructs can provide a selectable or screenable trait upon transcriptional activation (or inactivation) in response to a signal transduction pathway coupled to the target receptor. The indicator gene may be an unmodified gene already in the host cell pathway, such as the genes responsible for growth arrest in yeast. In certain embodiments a host cell gene may be operably linked to a "receptor-  
15 responsive" promoter. Alternatively, it may be a heterologous gene that has been so linked. Suitable genes and promoters are discussed below.

- To achieve optimal selection or screening, the host cell phenotype will be considered. For example, introducing a pheromone-responsive chimeric HIS3 gene into a yeast that has a wild-type HIS3 gene would frustrate genetic selection. Thus, to  
20 achieve nutritional selection, an auxotrophic strain is preferred. Yeast strains that are auxotrophic for histidine (HIS3) are known, see Struhl and Hill, (1987) *Mol. Cell. Biol.*, 7:104; Fasullo and Davis, *Mol. Cell. Biol.*, (1988) 8:4370. The HIS3 (imidazoleglycerol phosphate dehydratase) gene has been used as a selective marker in yeast. See Sikorski and Heiter, (1989) *Genetics*, 122:19; Struhl, et al., *P.N.A.S.* (1979) 76:1035; and, for  
25 FUS1-HIS3 fusions, see Stevenson, et al., (1992) *Genes Dev.*, 6:1293.

- In certain embodiments, the host cell can be modified in other ways. For example, when the host cell is a yeast cell it may be desirable to inactivate, such as by mutation or deletion, a homologous receptor, e.g., a pheromone receptor, present in the cell in order to minimize interference with signaling via the heterologous receptor.  
30 "Inactivation", with respect to genes of the host cell, means that production of a functional gene product is prevented or inhibited. Inactivation may be achieved by deletion of the gene, mutation of the promoter so that expression does not occur, or mutation of the coding sequence so that the gene product is inactive. Inactivation may be partial or total.

- 35 In a preferred embodiment of the subject assay, the yeast cells possess one or more of the following characteristics: (a) the endogenous FUS1 gene has been inactivated; (b) the endogenous SST2 gene, and/or other genes involved in desensitization, have been inactivated; (c) if there is a homologous, endogenous receptor gene it has been inactivated; and (d) if the yeast produces an endogenous ligand  
40 to the exogenous receptor, the genes encoding for the ligand been inactivated.

5 It is desirable that the exogenous receptor be exposed on a continuing basis to the peptides. In some instances, this may result in desensitization of the pheromone pathway to the stimulus. For example, the mating signal transduction pathway is known to become desensitized by several mechanisms including pheromone degradation and modification of the function of the receptor, G proteins and/or downstream elements of  
10 the pheromone signal transduction by the products of the SST2, STE50, AFR1 (Konopka, J.B. (1993) *Mol. Cell. Biol.* 13:6876-6888) and SGV1, MSG5, and SIG1 genes. Selected mutations in these genes can lead to hypersensitivity to pheromone and an inability to adapt to the presence of pheromone. For example, introduction of mutations that interfere with function into strains expressing heterologous G protein-  
15 coupled receptors constitutes a significant improvement on wild type strains and enables the development of extremely sensitive bioassays for compounds that interact with the receptors. Other mutations e.g. STE50, sgv1, bar1, ste2, ste3, pik1, msg5, sig1, and aft1, have the similar effect of increasing the sensitivity of the bioassay. Thus desensitization may be avoided by mutating (which may include deleting) the SST2 gene so that it no  
20 longer produces a functional protein, or by mutating one of the other genes listed above.

In certain embodiments, it will be desirable to complement the host yeast cells, e.g., at least partial function of an inactivated gene of the host cell can be supplied by an exogenous nucleic acid. For instance, yeast cells can be "mammalianized", and even "humanized", by complementation of receptor and signal transduction proteins with  
25 mammalian homologues. To illustrate, inactivation of a yeast Byr2/Stell1 gene can be complemented by expression of a human MEKK gene.

Complementations for use in the subject assay can be constructed without any undue experimentation. Indeed, many yeast genetic complementations with mammalian signal transduction proteins have been described in the art. For example, Mosteller et al.  
30 (1994) *Mol Cell Biol* 14:1104-12 demonstrates that human Ras proteins can complement loss of ras mutations in *S. cerevisiae*. Moreover, Toda et al. (1986) Princess Takamatsu Symp 17: 253-60 have shown that human ras proteins can complement the loss of RAS1 and RAS2 proteins in yeast, and hence are functionally homologous. Both human and yeast RAS proteins can stimulate the magnesium and  
35 guanine nucleotide-dependent adenylate cyclase activity present in yeast membranes. Ballester et al. (1989) *Cell* 59: 681-6 describe a vector to express the mammalian GAP protein in the yeast *S. cerevisiae*. When expressed in yeast, GAP inhibits the function of the human ras protein, and complements the loss of IRA1. IRA1 is a yeast gene that encodes a protein with homology to GAP and acts upstream of RAS. Mammalian GAP  
40 can therefore function in yeast and interact with yeast RAS. Wei et al. (1994) *Gene* 151: 279-84 describes that a human Ras-specific guanine nucleotide-exchange factor, Cdc25GEF, can complement the loss of CDC25 function in *S. cerevisiae*. Martegani et

5 al. (1992) EMBO J 11: 2151-7 describe the cloning by functional complementation of a mouse cDNA encoding a homolog of CDC25, a *Saccharomyces cerevisiae* RAS activator. Vojtek et al. (1993) J Cell Sci 105: 777-85 and Matviw et al. (1992) Mol Cell Biol 12: 5033-40 describe how a mouse CAP protein, e.g., an adenylyl cyclase associated protein associated with ras-mediated signal transduction, can complements  
10 defects in *S. cerevisiae*. Papasavvas et al. (1992) Biochem Biophys Res Commun 184:1378-85 also suggest that inactivated yeast adenylyl cyclase can be complemented by a mammalian adenylyl cyclase gene. Hughes et al. (1993) Nature 364: 349-52 describe the complementation of *byr1* in fission yeast by mammalian MAP kinase kinase (MEK). Parissenti et al. (1993) Mol Cell Endocrinol 98: 9-16 describes the reconstitution of  
15 bovine protein kinase C (PKC) in yeast. The Ca(2+) and phospholipid-dependent Ser/Thr kinase PKC plays important roles in the transduction of cellular signals in mammalian cells. Marcus et al. (1995) PNAS 92: 6180-4 suggests the complementation of *shk1* null mutations in *S. pombe* by either the structurally related *S. cerevisiae* Ste20 or mammalian p65PAK protein kinases.

20

## XII. Pharmaceutical Preparations of Identified Compounds

After identifying certain test compounds as potential surrogate ligands, or receptor antagonists, the practitioner of the subject assay will continue to test the efficacy and specificity of the selected compounds both *in vitro* and *in vivo*. Whether for  
25 subsequent *in vivo* testing, or for administration to an animal as an approved drug, agents identified in the subject assay can be formulated in pharmaceutical preparations for *in vivo* administration to an animal, preferably a human.

The compounds selected in the subject assay, or a pharmaceutically acceptable salt thereof, may accordingly be formulated for administration with a biologically  
30 acceptable medium, such as water, buffered saline, polyol (for example, glycerol, propylene glycol, liquid polyethylene glycol and the like) or suitable mixtures thereof. The optimum concentration of the active ingredient(s) in the chosen medium can be determined empirically, according to procedures well known to medicinal chemists. As used herein, "biologically acceptable medium" includes any and all solvents, dispersion  
35 media, and the like which may be appropriate for the desired route of administration of the pharmaceutical preparation. The use of such media for pharmaceutically active substances is known in the art. Except insofar as any conventional media or agent is incompatible with the activity of the compound, its use in the pharmaceutical preparation of the invention is  
40 contemplated. Suitable vehicles and their formulation inclusive of other proteins are described, for example, in the book *Remington's Pharmaceutical Sciences* (Remington's

5 Pharmaceutical Sciences. Mack Publishing Company, Easton, Pa., USA 1985). These vehicles include injectable "deposit formulations". Based on the above, such pharmaceutical formulations include, although not exclusively, solutions or freeze-dried powders of the compound in association with one or more pharmaceutically acceptable vehicles or diluents, and contained in buffered media at a suitable pH and isosmotic with  
10 physiological fluids. In preferred embodiment, the compound can be disposed in a sterile preparation for topical and/or systemic administration. In the case of freeze-dried preparations, supporting excipients such as, but not exclusively, mannitol or glycine may be used and appropriate buffered solutions of the desired volume will be provided so as to obtain adequate isotonic buffered solutions of the desired pH. Similar solutions may  
15 also be used for the pharmaceutical compositions of compounds in isotonic solutions of the desired volume and include, but not exclusively, the use of buffered saline solutions with phosphate or citrate at suitable concentrations so as to obtain at all times isotonic pharmaceutical preparations of the desired pH, (for example, neutral pH).

20 All patents, published patent applications and other references disclosed herein are hereby expressly incorporated by reference. In addition the contents of USSNs: 08/322,137; 08/463, 181; 08/655,192; 08/689,172; 08/936,632 and 08/946,298 are hereby incorporated by this reference.

25 **EXEMPLIFICATION**

The invention now being generally described will be more readily understood by reference to the following examples, which are included merely for purposes of illustration of certain aspects and embodiments of the present invention and are not  
30 intended to limit the invention.

**Example 1. Expression of the IL-8A Receptor in Cells**

35 The IL-8A receptor was expressed in both yeast and mammalian cells. For expression of the IL-8A receptor, an expression plasmid was derived from p1289, which contains the *LEU2* gene as a selectable marker and directs constitutive expression from the *PGK* promoter. The rabbit IL8A receptor was cloned from rabbit genomic DNA using the oligonucleotides "IL8R1FWD" (5'- CCCCCATGG  
40 AAGTAAACGTATGGAATATG-3') (SEQ ID NO:11 ) and "IL8R1REV" (5'- CCCTCTAGAGATTTGA AGGCACGTTGG-3') (SEQ ID NO:12). Cp2687 was constructed by subcloning the PCR product into p1289 using NcoI and XbaI as 5' and 3'

- 5 cloning sites. As the rabbit IL8AR gene contains an internal NcoI site, the receptor was, in fact, cloned as two pieces rejoined.

The yeast strain CY1141 (*MATa far1D1442 tbt1-1 fus1-HIS3 can1 ste14::trp1::LYS2 ste3D1156 gpa1(41)-Gai2 lys2 ura3 leu2 trp1 his3*) was constructed using standard genetic techniques and used for IL-8 receptor expression. The rabbit IL8A receptor couples to the pheromone response pathway in yeast through the Gpa1-Gai2 chimera.

For mammalian cell expression of the IL-8A receptor, HEK293 cells were used. HEK 293 cells stably expressing Ga16 were maintained in DMEM supplemented with 10% New Calf Serum and 200(g/ml G418). Cells at ~70% confluence were transfected with plasmids encoding the WT or mutant (made as described below) rabbit IL8A receptors and stable transfectants were selected in the presence of 400mg/ml hygromycin. Cells were maintained in selection medium for 7-9 days until the surviving cells expanded to 80-90% confluence.

## Example 2. Expression of Ligands of the IL-8 Receptor

There are two known human subtypes of the IL8 receptor. Subtype A (CXCR1) binds IL8 with high affinity ( $K_d = 0.1-1nM$ ) but shows very low affinity binding to GRO $\alpha$  and NAP-2 ( $K_d > 100nM$ ). Subtype B (CXCR2) binds all three ligands with high affinity.

Cells were constructed which coexpressed the IL-8 receptor and IL-8A receptor ligands. For ligand expression, plasmids were derived from the plasmid p1624 containing the *URA3* gene as a selectable marker and the *ADHI* promoter directing the constitutive expression of an Mfa signal sequence and leader. The ligand ORF was cloned into this plasmid using an AflII site at the 5' end and a BglII site at the 3' end to create a fusion junction of the indicated structure:

IL-8	LKR	SAKELRCQCI...	(SEQ ID NO: 13)
MGSA	LKR	ASVATELRCQCL...	(SEQ ID NO: 14)
40 NAP2	LKR	AELRCMCI...	(SEQ ID NO: 15)

The ligand genes were assembled by annealing the oligonucleotides described below with their complementary oligonucleotide, followed by ligation together into Cp1624 digested with AflII and BglII.

IL-8 (to yield the plasmid Cp1358):

IL8-1	TTAAGCGTGAGGCAGAAGCTTCTGCTAAGGAATTGAGATGTCAATGTATTAAGACTT (SEQ ID NO: 16)
IL8-2	ACTCTAAGCCATTCCATCCAAAGTTCAATTAAGGAATTGAGAGTTATTGAATCTGGTCCA (SEQ ID NO: 17)
50 IL8-3	CATTGTGCTAACACTGAAATTATTGTTAAGTTGTCTGATGGTAGAGAATTGTGTTTGGAT (SEQ ID NO: 18)
IL8-4	CCAAAGGAAAACCTGGGTTCAAAGAGTTGTTGAAAAGTTCTTGAAGAGAGCTGAAAACCTTGA (SEQ ID NO: 19)
IL8-5	TAGAGTAAGTCTTAATACATTGACATCTCAATTCCTTAGCAGAAGCTTCTGCCTCACGC (SEQ ID NO: 20)
IL8-6	ACAATGTGGACCAGATTCAATAACTCTCAATTCCTTAATGAACCTTGGATGGAATGGCT (SEQ ID NO: 21)

5 IL8-7 CTTTGGATCCAAACACAATTCTCTACCATCAGACAACTTAACAATAATTTTCAGTGTTAGC (SEQ ID NO: 22)  
IL8-8 GATCTCAAGAGTTTTTCAGCTCTCTTCAAGAACTTTTCAACAACCTTTGAACCCAGTTTTC (SEQ ID NO: 23)

The IL8 coding sequenced was subsequently amplified from Cp1358 using  
10 afl2-72-il8 (CCGCTTAAGCGTTCTGCTAAGGAATTGAGATGTC) (SEQ ID NO: 24)  
and PX25 (AAGTATATTGTATTTTGTACGAGC) (SEQ ID NO: 25), digested with  
AflII and BglII, and reintroduced into Cp1624 to create the junction indicated above.

15 MGSA/GRO $\alpha$  (to yield the plasmid Cp2685):

MGOPTF1 TTAAGCGTGCGTCCGTTGCTACAGAATTGAGGTGTCAATGTCTACAACTTTGCAA (SEQ ID NO: 26)  
MGOPTF2 GGTATCCACCCAAAGAACATTTCAGTCAGTTAAGTTAAGTCCCCAGGTCCACACTGTGC (SEQ ID NO: 27)  
MGOPTF3 TCAGACTGAAGTCATAGCTACATTAAAGAATGGTCGTAAGCCTGTTAAATCCTGC (SEQ ID NO: 28)  
20 MGOPTF4 ATCCCAATAGTAAAGAAAATCATCGAAAAGATGTTGAATAGTGATAAATCCAATTAA (SEQ ID NO: 29)  
MGOPTR1 GATCTTAATTGGATTATCACTATTCAACATCTTTTCGATGATTTTCTTACTATTGG (SEQ ID NO: 30)  
MGOPTR2 GGATGCAGGATTTAAACAGGCTTTACGACCATTCTTAATGTAGCTATGACTTCAGT (SEQ ID NO: 31)  
MGOPTR3 CTGAGCACAGTGTGGACCTGGGGACTTAACGTTAACTGACTGAATGTTCTTTGGGTGGA (SEQ ID NO: 32)  
25 MGOPTR4 TACCTTGCAAAGTTGTAGACATTGACACCTCAATTCTGTAGCAACGGACGCACGC (SEQ ID NO: 33)

CY9158 is the CY1141 strain harboring the plasmid (Cp2685) that expresses  
human GRO $\alpha$ . To achieve expression of secreted human GRO $\alpha$ , the coding sequence of  
GRO $\alpha$  was fused to sequences encoding the signal and leader sequences of the yeast  
30 *MFa1* gene, and placed this fusion protein under the control of the constitutive *ADHI*  
promoter.

NAP2 (to yield the plasmid Cp2674):

35 NAP2F1 TTAAGCGTGCTGAATTGAGATGTATGTGTATCAAGACCACCTCTGGTATCCACCC (SEQ ID NO: 34)  
NAP2F2 AAAGAACATCCAATCTTTGGAAGTTATCGGTAAGGGTACTCACTGTAACCAAGTT (SEQ ID NO: 35)  
NAP2F3 GAAGTTATCGCTACCTTGAAGGACGGTAGAAAGATTGTTTGGACCCAGACGCTC (SEQ ID NO: 36)  
NAP2F4 CAAGAATCAAGAAGATCGTTCAAAAGAAGTTGGCTGGTGACGAATCTGCTGACTA (SEQ ID NO: 37)  
40 NAP2R1 GATCTAGTCAGCAGATTTCGTCCAGCCAACTTCTTTGAACGATCTCTTGATT (SEQ ID NO: 38)  
NAP2R2 CTTGGAGCGTCTGGGTCCAAACAAATCTTTCTACCGTCCTTCAAGGTAGCGATAA (SEQ ID NO: 39)  
NAP2R3 CTTCAACTTGGTTACAGTGAGTACCCTTACCGATAACTTCCAAAGATTGGATGT (SEQ ID NO: 40)  
NAP2R4 CTTTGGGTGGATACCAGAGGTGGTCTTGATACACATACATCTCAATTCAGCACGC (SEQ ID NO: 41)

45

## Example 2. Creation of IL-8A Receptor Mutants

By random mutagenesis of the IL-8 receptor, mutants that show altered coupling  
50 behavior to the ligands GRO $\alpha$  and NAP-2, as well as showing an enhanced response to  
IL8 have been isolated.

The plasmid Cp2687, encoding the gene for the rabbit IL8A receptor under the  
control of the constitutive *PGK1* promoter, was randomly mutagenized using  
55 hydroxylamine and introduced into CY9158 (which expresses human GRO $\alpha$ ).  
Hydroxylamine mutagenesis was performed by incubating 10 mg Cp2687 (containing



5 the IL-8A receptor) in 500 ml of 7% HONH<sub>2</sub>•HCl. 1.8% NaOH for 24 hours at 37°. The DNA was ethanol precipitated, resuspended in ddH<sub>2</sub>O and introduced into CY9158 using lithium acetate.

10 Approximately 30,000 yeast colonies were selected on SCLU (SCLU contains a mixture of amino acids lacking leucine and uracil; the pH was adjusted to 6.8 using 4.5 ml 1M KOH and 25 ml 1M K-Pipes pH6.8 per liter), and after 2 days of growth at 30°C, the colonies were replica plated to the same media lacking histidine and containing 1mM 3-aminotriazole (AT, a competitive inhibitor of the His3 gene product) (SCH1 media). Yeast cells expressing the wild-type receptor do not grow on this media  
15 since GRO $\alpha$  activates this receptor poorly if at all; the *fus1-HIS3* construct, which serves as a measure of pathway activation, is therefore repressed and the cells remain auxotrophic for histidine. Therefore, only cells expressing receptors that have gained the ability to be activated by GRO $\alpha$  are selected on this media. Approximately 200 such colonies were identified. Figure 1 shows the GRO $\alpha$  and NAP-2 dependent growth of  
20 yeast strains expressing the mutant receptors, as well as the inability of these ligands to activate wild-type receptor.

25 Sequencing of the receptor plasmids showed that all the mutations found were in the intracellular loops and the C-terminal tail (Fig.2). Using single amino acid code, the mutations identified were:

R73W (i.e., at position 73 an Arginine to Tryptophan mutation) – (CGG to TGG)  
M246I – (ATG to ATA)  
G320R – (GGA to AGA)

30

Subsequent segregation of Cp2685 (*URA3*<sup>+</sup> *PGKp*-GRO $\alpha$ ) demonstrated that the His<sup>+</sup> phenotype was dependent on the presence of GRO $\alpha$ : Ura<sup>-</sup> progeny selected by their ability to grow in the presence of 5-fluoro orotic acid were no longer able to grow  
35 on SCLH1. Plasmid DNA was recovered from colonies that showed GRO $\alpha$ -dependent growth and reintroduced into CY9158 to ensure that the mutated DNA was able to confer the identical His<sup>+</sup> phenotype to a naïve host.

40 Plasmids expressing double mutants were constructed by replacing a BamHI/XbaI fragment from the R73W mutant with that from either the M246I or the G320R mutants. When these mutations are combined, the growth phenotype is enhanced. This is clear when their growth is compared on media containing 10mM 3-AT, a concentration which demands higher expression of His3 to confer histidine prototrophy (Fig.3).

45

$\beta$ -galactosidase activity arising from the expression of a *fus1-lacZ* plasmid (Cp1584) can be measured either in autocrine strains (expressing ligand endogenously) or in cells that are supplied with agonist exogenously. In autocrine strains, the response to IL-8 and NAP-2 is consistent with the results of the *fus1-HIS3* assay, but the response  
50 of these mutants to GRO $\alpha$  is greatly diminished (Fig. 4). The reason for this discrepancy may lie in the fact that the growth assay measures activation over a period of days, while

5 the *fus1-lacZ* assay measures accumulation of enzyme over a period of a few hours. The  
b-galactosidase assays were performed using ONPG as substrate (Mol. Cell. Biol.  
7:2316). For CPRG assays, 100ml of cells in synthetic media were mixed with 20ml of  
1.6 mg/ml CPRG in 2.5% Triton X-100/138 mM K-Pipes pH6.8, and incubated at 37°C.  
10 The reactions were stopped by the addition of 10ml of 1M Na<sub>2</sub>CO<sub>3</sub> and the absorbance  
was measured at 578 nm. Dose responses using IL8, GRO $\alpha$  or NAP-2 (R&D Systems)  
were performed by incubating logarithmic cultures in SCLUT (SCLUT lacks leucine,  
uracil and tryptophan).with the indicated amount of agonist for the indicated times at 30°  
prior to the addition of the lysis/substrate buffer.

15 The results of such assays with IL8, GRO $\alpha$  or NAP-2 being provided  
exogenously are shown in Figure 5. When ligand is applied exogenously, *fus1-lacZ*  
activity in yeast strains expressing the rabbit IL8AR increases in a dose dependent  
manner. The response of all the single mutants and at least one of the double mutants  
20 (R73W M246I) to IL8 is significantly increased with respect to wild type receptor (Fig.  
5A). In the assay with NAP-2, both double mutants and the M246I mutant show  
enhanced coupling over the wild type receptor (Fig. 5C). EC<sub>50</sub> values could not be  
obtained since the assay was not saturated at the highest ligand concentration (5M), an  
obstacle that is magnified in the GRO $\alpha$  assay (Fig. 5B).

25 Ligand binding assays with membranes from yeast strains expressing the WT  
rIL8 receptor or the double mutant R73W G320R, showed a slight difference in binding  
affinity for either IL8 or GRO( (Fig. 6), although the mutant showed a 2-fold increase in  
the B<sub>max</sub>.

30 In order to study the effects of these mutations on receptor function in  
mammalian cells, the mutant receptors were cloned into a mammalian expression vector  
and transfected into HEK293/Ga16 cells, which stably express a Ga16 construct. Ca<sup>++</sup>  
fluxes in response to agonist were measured in a dose response assay and in a pertussis  
35 toxin experiment. To measure Ca<sup>++</sup> flux, transfected cells were transferred to black-  
walled 96-well plates at 9x10<sup>6</sup> cells/well. After 24 hours of growth in the presence or  
absence of pertussis toxin (0.5 mg per ml; Calbiochem), the growth medium was  
removed and the cells were loaded with the fluorescent dye Fluo-3 for 1 hr at 37° in the  
presence of 20mM Probenicid (Sigma). Each well was washed with Hanks buffered salt  
40 solution/20mM HEPES/20mM Probenicid. The appropriate amount of agonist was  
added and fluorescence intensity was measured over a period of 5 minutes using a  
Fluorescence Imaging Plate Reader (Molecular Devices).

45 The dose response (Fig. 8) indicates that the M246I mutant recapitulates the  
novel coupling behavior seen in yeast cells, in that it displays a leftward shift in the  
MGSA dose response curve, suggesting a higher apparent potency of that agonist on the  
mutant receptor as compared to wild type. In addition, this mutation in conjunction with  
the R73W mutation confers a marked sensitivity to pertussis toxin inhibition for IL-8  
and to a lesser extent for MGSA/GRO $\alpha$ , but not for NAP2 (Fig. 9). These data suggest  
50 that these mutations alter the coupling of the IL8 receptor to the G-protein. The  
mutations may facilitate the ability of a heterologous G-protein to recognize and undergo

5 guanine nucleotide exchange in yeast. Alternatively, it is possible that some of these mutations have affected the binding affinities of the receptor with respect to the various ligands.

### 10 **Example 3. Creation of Mutants of the Galanin Receptor-1**

One of the mutations identified in the IL-8 receptor has been transferred to human galanin receptor-1 to enhance signaling by that molecule through Gpa1-Gαi2.

15 A DNA fragment encoding the human galanin receptor-1 (hGalr1), generated by the PCR primers GLNRPCRf (5'-GTGACTGGTCTGCCATGGAGCTGGCGGTCGGGAACCT-3') (SEQ ID NO: 42) and GLNRPCRR (5'-CGCGGATCCCACATGAGTACAATTGGT-3') (SEQ ID NO: 43) from a cloned template (pCR2-hGalr; E. Davis, BMS) was introduced into the yeast expression vector Cp4258 (*LEU2* selectable marker, *PGK* promoter, *MFal* signal and leader sequences) using *NcoI* and *XbaI*. The A(R mutation was introduced using the Quik-Change mutagenesis protocol (Stratagene) with the oligonucleotide PX1267 (5' - CTGAAAATT TCAGGAAGAGATATAACAAGTG TTCAAG - 3') (SEQ ID NO: 44) and PX1268 (5' - CTTGAACA CTTGTTTATATCTCTTCCTGAAA TTTTCAG) (SEQ ID NO: 45).

Oligonucleotides used for PCR cloning from yeast expression vectors were R1CEP4fwd (5'-CCCAAGCT TGCCACCATGGAAGTAAACGTATG-3') (SEQ ID NO: 46) and R1CEP4rev (5'-CCCCTCGAGCTAGAGATTTGAAGGCACGTT-3') (SEQ ID NO: 47). The PCR products were cut with *HindIII* and *XhoI* and cloned into pCEP4 (Invitrogen) cut with the same enzymes. The clones were sequenced prior to use.

35 The wild type human galanin receptor-1 couples somewhat weakly to the pheromone response pathway in yeast. In order to improve this response, the amino acid sequences of the rabbit IL8 receptor and the human galanin receptor-1 were compared a region of limited homology in the seventh transmembrane domain and the C-terminal tail was identified. The galanin receptor contains an Ala in the same position as Gly320 of the rabbit IL8 receptor:

LGFLHSCLNPIIYAFIGQNFRNGFLKM rIL8aR (SEQ ID NO: 3)  
LAYSNSSVNPIIYAFLSENFRKAYKQV hGalR1 (SEQ ID NO: 5)

By changing the Ala (underlined above) to an Arg, the galanin receptor showed an enhanced response to ligand (Fig.7).

50

5

EQUIVALENTS

10 Those skilled in the art will recognize, or be able to ascertain using no more than routine experimentation, many equivalents to the specific embodiments of the invention described herein. Such equivalents are intended to be encompassed by the following claims.

11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65  
66  
67  
68  
69  
70  
71  
72  
73  
74  
75  
76  
77  
78  
79  
80  
81  
82  
83  
84  
85  
86  
87  
88  
89  
90  
91  
92  
93  
94  
95  
96  
97  
98  
99  
100  
101  
102  
103  
104  
105  
106  
107  
108  
109  
110  
111  
112  
113  
114  
115  
116  
117  
118  
119  
120  
121  
122  
123  
124  
125  
126  
127  
128  
129  
130  
131  
132  
133  
134  
135  
136  
137  
138  
139  
140  
141  
142  
143  
144  
145  
146  
147  
148  
149  
150  
151  
152  
153  
154  
155  
156  
157  
158  
159  
160  
161  
162  
163  
164  
165  
166  
167  
168  
169  
170  
171  
172  
173  
174  
175  
176  
177  
178  
179  
180  
181  
182  
183  
184  
185  
186  
187  
188  
189  
190  
191  
192  
193  
194  
195  
196  
197  
198  
199  
200  
201  
202  
203  
204  
205  
206  
207  
208  
209  
210  
211  
212  
213  
214  
215  
216  
217  
218  
219  
220  
221  
222  
223  
224  
225  
226  
227  
228  
229  
230  
231  
232  
233  
234  
235  
236  
237  
238  
239  
240  
241  
242  
243  
244  
245  
246  
247  
248  
249  
250  
251  
252  
253  
254  
255  
256  
257  
258  
259  
260  
261  
262  
263  
264  
265  
266  
267  
268  
269  
270  
271  
272  
273  
274  
275  
276  
277  
278  
279  
280  
281  
282  
283  
284  
285  
286  
287  
288  
289  
290  
291  
292  
293  
294  
295  
296  
297  
298  
299  
300  
301  
302  
303  
304  
305  
306  
307  
308  
309  
310  
311  
312  
313  
314  
315  
316  
317  
318  
319  
320  
321  
322  
323  
324  
325  
326  
327  
328  
329  
330  
331  
332  
333  
334  
335  
336  
337  
338  
339  
340  
341  
342  
343  
344  
345  
346  
347  
348  
349  
350  
351  
352  
353  
354  
355  
356  
357  
358  
359  
360  
361  
362  
363  
364  
365  
366  
367  
368  
369  
370  
371  
372  
373  
374  
375  
376  
377  
378  
379  
380  
381  
382  
383  
384  
385  
386  
387  
388  
389  
390  
391  
392  
393  
394  
395  
396  
397  
398  
399  
400  
401  
402  
403  
404  
405  
406  
407  
408  
409  
410  
411  
412  
413  
414  
415  
416  
417  
418  
419  
420  
421  
422  
423  
424  
425  
426  
427  
428  
429  
430  
431  
432  
433  
434  
435  
436  
437  
438  
439  
440  
441  
442  
443  
444  
445  
446  
447  
448  
449  
450  
451  
452  
453  
454  
455  
456  
457  
458  
459  
460  
461  
462  
463  
464  
465  
466  
467  
468  
469  
470  
471  
472  
473  
474  
475  
476  
477  
478  
479  
480  
481  
482  
483  
484  
485  
486  
487  
488  
489  
490  
491  
492  
493  
494  
495  
496  
497  
498  
499  
500  
501  
502  
503  
504  
505  
506  
507  
508  
509  
510  
511  
512  
513  
514  
515  
516  
517  
518  
519  
520  
521  
522  
523  
524  
525  
526  
527  
528  
529  
530  
531  
532  
533  
534  
535  
536  
537  
538  
539  
540  
541  
542  
543  
544  
545  
546  
547  
548  
549  
550  
551  
552  
553  
554  
555  
556  
557  
558  
559  
560  
561  
562  
563  
564  
565  
566  
567  
568  
569  
570  
571  
572  
573  
574  
575  
576  
577  
578  
579  
580  
581  
582  
583  
584  
585  
586  
587  
588  
589  
590  
591  
592  
593  
594  
595  
596  
597  
598  
599  
600  
601  
602  
603  
604  
605  
606  
607  
608  
609  
610  
611  
612  
613  
614  
615  
616  
617  
618  
619  
620  
621  
622  
623  
624  
625  
626  
627  
628  
629  
630  
631  
632  
633  
634  
635  
636  
637  
638  
639  
640  
641  
642  
643  
644  
645  
646  
647  
648  
649  
650  
651  
652  
653  
654  
655  
656  
657  
658  
659  
660  
661  
662  
663  
664  
665  
666  
667  
668  
669  
670  
671  
672  
673  
674  
675  
676  
677  
678  
679  
680  
681  
682  
683  
684  
685  
686  
687  
688  
689  
690  
691  
692  
693  
694  
695  
696  
697  
698  
699  
700  
701  
702  
703  
704  
705  
706  
707  
708  
709  
710  
711  
712  
713  
714  
715  
716  
717  
718  
719  
720  
721  
722  
723  
724  
725  
726  
727  
728  
729  
730  
731  
732  
733  
734  
735  
736  
737  
738  
739  
740  
741  
742  
743  
744  
745  
746  
747  
748  
749  
750  
751  
752  
753  
754  
755  
756  
757  
758  
759  
760  
761  
762  
763  
764  
765  
766  
767  
768  
769  
770  
771  
772  
773  
774  
775  
776  
777  
778  
779  
780  
781  
782  
783  
784  
785  
786  
787  
788  
789  
790  
791  
792  
793  
794  
795  
796  
797  
798  
799  
800  
801  
802  
803  
804  
805  
806  
807  
808  
809  
810  
811  
812  
813  
814  
815  
816  
817  
818  
819  
820  
821  
822  
823  
824  
825  
826  
827  
828  
829  
830  
831  
832  
833  
834  
835  
836  
837  
838  
839  
840  
841  
842  
843  
844  
845  
846  
847  
848  
849  
850  
851  
852  
853  
854  
855  
856  
857  
858  
859  
860  
861  
862  
863  
864  
865  
866  
867  
868  
869  
870  
871  
872  
873  
874  
875  
876  
877  
878  
879  
880  
881  
882  
883  
884  
885  
886  
887  
888  
889  
890  
891  
892  
893  
894  
895  
896  
897  
898  
899  
900  
901  
902  
903  
904  
905  
906  
907  
908  
909  
910  
911  
912  
913  
914  
915  
916  
917  
918  
919  
920  
921  
922  
923  
924  
925  
926  
927  
928  
929  
930  
931  
932  
933  
934  
935  
936  
937  
938  
939  
940  
941  
942  
943  
944  
945  
946  
947  
948  
949  
950  
951  
952  
953  
954  
955  
956  
957  
958  
959  
960  
961  
962  
963  
964  
965  
966  
967  
968  
969  
970  
971  
972  
973  
974  
975  
976  
977  
978  
979  
980  
981  
982  
983  
984  
985  
986  
987  
988  
989  
990  
991  
992  
993  
994  
995  
996  
997  
998  
999  
1000  
1001  
1002  
1003  
1004  
1005  
1006  
1007  
1008  
1009  
1010  
1011  
1012  
1013  
1014  
1015  
1016  
1017  
1018  
1019  
1020  
1021  
1022  
1023  
1024  
1025  
1026  
1027  
1028  
1029  
1030  
1031  
1032  
1033  
1034  
1035  
1036  
1037  
1038  
1039  
1040  
1041  
1042  
1043  
1044  
1045  
1046  
1047  
1048  
1049  
1050  
1051  
1052  
1053  
1054  
1055  
1056  
1057  
1058  
1059  
1060  
1061  
1062  
1063  
1064  
1065  
1066  
1067  
1068  
1069  
1070  
1071  
1072  
1073  
1074  
1075  
1076  
1077  
1078  
1079  
1080  
1081  
1082  
1083  
1084  
1085  
1086  
1087  
1088  
1089  
1090  
1091  
1092  
1093  
1094  
1095  
1096  
1097  
1098  
1099  
1100  
1101  
1102  
1103  
1104  
1105  
1106  
1107  
1108  
1109  
1110  
1111  
1112  
1113  
1114  
1115  
1116  
1117  
1118  
1119  
1120  
1121  
1122  
1123  
1124  
1125  
1126  
1127  
1128  
1129  
1130  
1131  
1132  
1133  
1134  
1135  
1136  
1137  
1138  
1139  
1140  
1141  
1142  
1143  
1144  
1145  
1146  
1147  
1148  
1149  
1150  
1151  
1152  
1153  
1154  
1155  
1156  
1157  
1158  
1159  
1160  
1161  
1162  
1163  
1164  
1165  
1166  
1167  
1168  
1169  
1170  
1171  
1172  
1173  
1174  
1175  
1176  
1177  
1178  
1179  
1180  
1181  
1182  
1183  
1184  
1185  
1186  
1187  
1188  
1189  
1190  
1191  
1192  
1193  
1194  
1195  
1196  
1197  
1198  
1199  
1200  
1201  
1202  
1203  
1204  
1205  
1206  
1207  
1208  
1209  
1210  
1211  
1212  
1213  
1214  
1215  
1216  
1217  
1218  
1219  
1220  
1221  
1222  
1223  
1224  
1225  
1226  
1227  
1228  
1229  
1230  
1231  
1232  
1233  
1234  
1235  
1236  
1237  
1238  
1239  
1240  
1241  
1242  
1243  
1244  
1245  
1246  
1247  
1248  
1249  
1250  
1251  
1252  
1253  
1254  
1255  
1256  
1257  
1258  
1259  
1260  
1261  
1262  
1263  
1264  
1265  
1266  
1267  
1268  
1269  
1270  
1271  
1272  
1273  
1274  
1275  
1276  
1277  
1278  
1279  
1280  
1281  
1282  
1283  
1284  
1285  
1286  
1287  
1288  
1289  
1290  
1291  
1292  
1293  
1294  
1295  
1296  
1297  
1298  
1299  
1300  
1301  
1302  
1303  
1304  
1305  
1306  
1307  
1308  
1309  
1310  
1311  
1312  
1313  
1314  
1315  
1316  
1317  
1318  
1319  
1320  
1321  
1322  
1323  
1324  
1325  
1326  
1327  
1328  
1329  
1330  
1331  
1332  
1333  
1334  
1335  
1336  
1337  
1338  
1339  
1340  
1341  
1342  
1343  
1344  
1345  
1346  
1347  
1348  
1349  
1350  
1351  
1352  
1353  
1354  
1355  
1356  
1357  
1358  
1359  
1360  
1361  
1362  
1363  
1364  
1365  
1366  
1367  
1368  
1369  
1370  
1371  
1372  
1373  
1374  
1375  
1376  
1377  
1378  
1379  
1380  
1381  
1382  
1383  
1384  
1385  
1386  
1387  
1388  
1389  
1390  
1391  
1392  
1393  
1394  
1395  
1396  
1397  
1398  
1399  
1400  
1401  
1402  
1403  
1404  
1405  
1406  
1407  
1408  
1409  
1410  
1411  
1412  
1413  
1414  
1415  
1416  
1417  
1418  
1419  
1420  
1421  
1422  
1423  
1424  
1425  
1426  
1427  
1428  
1429  
1430  
1431  
1432  
1433  
1434  
1435  
1436  
1437  
1438  
1439  
1440  
1441  
1442  
1443  
1444  
1445  
1446  
1447  
1448  
1449  
1450  
1451  
1452  
1453  
1454  
1455  
1456  
1457  
1458  
1459  
1460  
1461  
1462  
1463  
1464  
1465  
1466  
1467  
1468  
1469  
1470  
1471  
1472  
1473  
1474  
1475  
1476  
1477  
1478  
1479  
1480  
1481  
1482  
1483  
1484  
1485  
1486  
1487  
1488  
1489  
1490  
1491  
1492  
1493  
1494  
1495  
1496  
1497  
1498  
1499  
1500  
1501  
1502  
1503  
1504  
1505  
1506  
1507  
1508  
1509  
1510  
1511  
1512  
1513  
1514  
1515  
1516  
1517  
1518  
1519  
1520  
1521  
1522  
1523  
1524  
1525  
1526  
1527  
1528  
1529  
1530  
1531  
1532  
1533  
1534  
1535  
1536  
1537  
1538  
1539  
1540  
1541  
1542  
1543  
1544  
1545  
1546  
1547  
1548  
1549  
1550  
1551  
1552  
1553  
1554  
1555  
1556  
1557  
1558  
1559  
1560  
1561  
1562  
1563  
1564  
1565  
1566  
1567  
1568  
1569  
1570  
1571  
1572  
1573  
1574  
1575  
1576  
1577  
1578  
1579  
1580  
1581  
1582  
1583  
1584  
1585  
1586  
1587  
1588  
1589  
1590  
1591  
1592  
1593  
1594  
1595  
1596  
1597  
1598  
1599  
1600  
1601  
1602  
1603  
1604  
1605  
1606  
1607  
1608  
1609  
1610  
1611  
1612  
1613  
1614  
1615  
1616  
1617  
1618  
1619  
1620  
1621  
1622  
1623  
1624  
1625  
1626  
1627  
1628  
1629  
1630  
1631  
1632  
1633  
1634  
1635  
1636  
1637  
1638  
1639  
1640  
1641  
1642  
1643  
1644  
1645  
1646  
1647  
1648  
1649  
1650  
1651  
1652  
1653  
1654  
1655  
1656  
1657  
1658  
1659  
1660  
1661  
1662  
1663  
1664  
1665  
1666  
1667  
1668  
1669  
1670  
1671  
1672  
1673  
1674  
1675  
1676  
1677  
1678  
1679  
1680  
1681  
1682  
1683  
1684  
1685  
1686  
1687  
1688  
1689  
1690  
1691  
1692  
1693  
1694  
1695  
1696  
1697  
1698  
1699  
1700  
1701  
1702  
1703  
1704  
1705  
1706  
1707  
1708  
1709  
1710  
1711  
1712  
1713  
1714  
1715  
1716  
1717  
1718  
1719  
1720  
1721  
1722  
1723  
1724  
1725  
1726  
1727  
1728  
1729  
1730  
1731  
1732  
1733  
1734  
1735  
1736  
1737  
1738  
1739  
1740  
1741  
1742  
1743  
1744  
1745  
1746  
1747  
1748  
1749  
1750  
1751  
1752  
1753  
1754  
1755  
1756  
1757  
1758  
1759  
1760  
1761  
1762  
1763  
1764  
1765  
1766  
1767  
1768  
1769  
1770  
1771  
1772  
1773  
1774  
1775  
1776  
1777  
1778  
1779  
1780  
1781  
1782  
1783  
1784  
1785  
1786  
1787  
1788  
1789  
1790  
1791  
1792  
1793  
1794  
1795  
1796  
1797  
1798  
1799  
1800  
1801  
1802  
1803  
1804  
1805  
1806  
1807  
1808  
1809  
1810  
1811  
1812  
1813  
1814  
1815  
1816  
1817  
1818  
1819  
1820  
1821  
1822  
1823  
1824  
1825  
1826  
1827  
1828  
1829  
1830  
1831  
1832  
1833  
1834  
1835  
1836  
1837  
1838  
1839  
1840  
1841  
1842  
1843  
1844  
1845  
1846  
1847  
1848  
1849  
1850  
1851  
1852  
1853  
1854  
1855  
1856  
1857  
1858  
1859  
1860  
1861  
1862  
1863  
1864  
1865  
1866  
1867  
1868  
1869  
1870  
1871  
1872  
1873  
1874  
1875  
1876  
1877  
1878  
1879  
1880  
1881  
1882  
1883  
1884  
1885  
1886  
1887  
1888  
1889  
1890  
1891  
1892  
1893  
1894  
1895  
1896  
1897  
1898  
1899  
1900  
1901  
1902  
1903  
1904  
1905  
1906  
1907  
1908  
1909  
1910  
1911  
1912  
1913  
1914  
1915  
1916  
1917  
1918  
1919  
1920  
1921  
1922  
1923  
1924  
1925  
1926  
1927  
1928  
1929  
1930  
1931  
1932  
1933  
1934  
1935  
1936  
1937  
1938  
1939  
1940  
1941  
1942  
1943  
1944  
1945  
1946  
1947  
1948  
1949  
1950  
1951  
1952  
1953  
1954  
1955  
1956  
1957  
1958  
1959  
1960  
1961  
1962  
1963  
1964  
1965  
1966  
1967  
1968  
1969  
1970  
1971  
1972  
1973  
1974  
1975  
1976  
1977  
1978  
1979  
1980  
1981  
1982  
1983  
1984  
1985  
1986  
1987  
1988  
1989  
1990  
1991  
1992  
1993  
1994  
1995  
1996  
1997  
1998  
1999  
2000  
2001  
2002  
2003  
2004  
2005  
2006  
2007  
2008  
2009  
2010  
2011  
2012  
2013  
2014  
2015  
2016  
2017  
2018  
2019  
2020  
2021  
2022  
2023  
2024  
2025  
2026  
2027  
2028  
2029  
2030  
2031  
2032  
2033  
2034  
2035  
2036  
2037  
2038  
2039  
2040  
2041  
2042  
2043  
2044  
2045  
2046  
2047  
2048  
2049  
2050  
2051  
2052  
2053  
2054  
2055  
2056  
2057  
2058  
2059  
2060  
2061  
2062  
2063  
2064  
2065  
2066  
2067  
2068  
2069  
2070  
2071  
2072  
2073  
2074  
2075  
2076  
2077  
2078  
2079  
2080  
2081  
2082  
2083  
2084  
2085  
2086  
2087  
2088  
2089  
2090  
2091  
2092  
2093  
2094  
2095  
2096  
2097  
2098  
2099  
2100  
2101  
2102  
2103  
2104  
2105  
2106  
2107  
2108  
2109  
2110  
2111  
2112  
2113  
2114  
2115  
2116  
2117  
2118  
2119  
2120  
2121  
2122  
2123  
2124  
2125  
2126  
2127  
2128  
2129  
2130  
2131  
2132  
2133  
2134  
2135  
2136  
2137  
2138  
2139  
2140  
2141  
2142  
2143  
2144  
2145  
2146  
2147  
2148  
2149  
2150  
2151  
2152  
2153  
2154  
2155  
2156  
2157  
2158  
2159  
2160  
2161  
2162  
2163  
2164  
2165  
2166  
2167  
2168  
2169  
2170  
2171  
2172  
2173  
2174  
2175  
2176  
2177  
2178  
2179  
2180  
2181  
2182  
2183  
2184  
2185  
2186  
2187  
2188  
2189  
2190  
2191  
2192  
2193  
2194  
2195  
2196  
2197  
2198  
2199  
2200  
2201  
2202  
2203  
2204  
2205  
2206  
2207  
2208  
2209  
2210  
2211  
2212  
2213  
2214  
2215  
2216  
2217  
2218